

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Claudia BERGER et al  
Serial No.: To Be Determined      Group Art Unit: To Be Determined  
Filed: November 18, 2003      Examiner: To Be Determined  
Title: COMPOUNDS WITH NEP/MP-INHIBITORY ACTIVITY AND  
USES THEREOF

CLAIM FOR PRIORITY UNDER 35 U.S.C. §119


Mail Stop Patent Application  
Director of USPTO  
P.O. Box 1450  
Alexandria, VA 22313-1450

Sir:

The benefit of the filing date of prior foreign application No. 01 11  
2231.4, filed in the European Patent Office on May 18, 2001, and U.S. patent  
application No. 60/292,337, filed May 22, 2001, is hereby requested and the right  
of priority under 35 U.S.C. §119 is hereby claimed.

In support of this claim, filed herewith is a certified copy of the original  
foreign application.

Respectfully submitted,




J. D. Evans  
Registration No. 26,269  
Christopher T. McWhinney  
Registration No. 42,875

Date: November 18, 2003

CROWELL & MORING, LLP  
P.O. Box 14300  
Washington, DC 20044-4300  
Telephone No.: (202) 624-2500  
Facsimile No.: (202) 628-8844

JDE/CTM/lw





**Eur päisches  
Patentamt**

**European  
Patent Office**

**Office européen  
des brevets**

**Bescheinigung**

**Certificate**

**Attestation**

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

**Patentanmeldung Nr.    Patent application No.    Demande de brevet n°**

01112231.4

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

**R C van Dijk**

DEN HAAG, DEN  
THE HAGUE,    08/08/02  
LA HAYE, LE





Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

**Blatt 2 der Bescheinigung**  
**Sheet 2 of the certificate**  
**Page 2 de l'attestation**

Anmeldung Nr.:  
Application no.: 01112231.4  
Demande n°:

Anmeldetag:  
Date of filing: 18/05/01  
Date de dépôt:

Anmelder:  
Applicant(s):  
Demandeur(s):  
Solvay Pharmaceuticals GmbH  
30173 Hannover  
GERMANY

Bezeichnung der Erfindung:  
Title of the invention:  
Titre de l'invention:

Use of compounds with combined nep/mp-inhibitory activity in the preparation of medicaments

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat:  
State:  
Pays:

Tag:  
Date:  
Date:

Aktenzeichen:  
File no.  
Numéro de dépôt:

Internationale Patentklassifikation:  
International Patent classification:  
Classification internationale des brevets:

A61K38/05, A61P9/00

Am Anmeldetag benannte Vertragsstaaten:  
Contracting states designated at date of filing:  
Etats contractants désignés lors du dépôt:

AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE/TR

Bemerkungen:  
Remarks:  
Remarques:





HA0117EP30

1

**Use of Compounds with Combined NEP/MP-Inhibitory Activity  
in the Preparation of Medicaments**

**Description****FIELD OF THE INVENTION**

5           This invention relates to the new medical use of compounds which act as combined or concurrent inhibitors of neutral endopeptidase (NEP) and of a specific metalloprotease (MP) which recently has been cloned and newly identified as a genuine metalloprotease with broad substrate specificity.

**10           BACKGROUND OF THE INVENTION**

          Metalloproteases are polypeptides which form a particular family of structurally and functionally related enzymes, e.g. peptidases, which are of pharmaceutical or pharmacological interest in the context of treatment or prophylaxis of various diseases. Several diseases have been identified where metalloproteases play a critical role in the pathology of the disease. For example,  
15           a number of zinc metalloproteases or particular families of structurally and functionally related enzymes have been identified and characterized in the state of the art, and it has become apparent that the participation of these enzymes, e.g. zinc metalloproteases, plays a role in a diverse array of biological functions encompassing both normal and disease situations. Zinc metalloproteases are subset of such enzymes whose catalytic functions are critically dependent on  
20           the zinc ion at the active site. This group of enzymes, which comprises various families classified on the basis of both sequence and structural information, are for example described to be intimately involved in such processes as e.g. embryonic development, cartilage and bone formation, processing of peptide hormones, reproduction, cardiovascular diseases, arthritis and cancer. Thus, there is particular interest in the pharmaceutical art not only to investigate the key  
25           roles of each of the metalloproteases and their potential interrelationship in health and disease, but especially also in designing improved therapeutical concepts for the management of diseases involving said metalloproteases.

          On the basis of sequence and structural information around the zinc binding-site of the  
30           zinc metalloproteases these enzymes may be classified into several families which may be further classified into superfamilies such as the "metzincins" (astacin, serratio, reprolysin, matrixin), the "gluzincins" (thermolysin, neprilysin, angiotensin converting enzyme, aminopeptidase), or the "zincins" comprising the superfamilies of metzincins and gluzincins. Such grouping not only aids in the elucidation of common catalytic and biosynthetic processing mechanisms, but also is  
35           invaluable in elucidating the function(s) of newly identified proteins which possess similar zinc

binding motifs. Some individual examples of metalloproteases, e.g. zinc enzymes, already identified in the state of the art comprise neprilysin, endothelin converting enzyme, angiotensin converting enzyme, thermolysin, aminopeptidase, astacin, serratin, reprotin, matrixin, insulinase, carboxypeptidase and DD-carboxypeptidase.

5

Some more specific features and related known activities of particularly interesting metalloprotease subtypes like neutral endopeptidase (NEP), endothelin converting enzyme (ECE), and angiotensin converting enzyme (ACE) may be summarized as follows.

10       Angiotensin I Converting Enzyme (ACE; peptidyl dipeptidase A; EC 3.4.15.1) is a member of the angiotensin converting enzyme family of zinc metalloproteases. ACE is primarily expressed at the surface of endothelial, epithelial and neuroepithelial cells (somatic ACE) as an ectoenzyme, meaning that it is anchored to the plasma membrane with the bulk of its mass, including its catalytic site/s, facing the extracellular milieu. ACE is found in the plasma membrane of vascular  
15       endothelial cells, with high levels found at the vascular endothelial surface of the lung such that the active sites of ACE are posed to metabolize circulating substrates. In addition to the endothelial location of ACE, the enzyme is also expressed in the brush borders of absorptive epithelia of the small intestine and the kidney proximal convoluted tubule. ACE is also found in mononuclear cells, such as monocytes after macrophage differentiation and T-lymphocytes, and in  
20       fibroblasts. In vitro autoradiography, employing radiolabelled specific ACE inhibitors, and immunohistochemical studies have mapped the principal locations of ACE in the brain. ACE was found primarily in the choroid plexus, which may be the source of ACE in cerebrospinal fluid, ependyma, subfornical organ, basal ganglia (caudate-putamen and globus pallidus), substantia nigra and pituitary. A soluble form of ACE has been detected in many biological fluids such as  
25       serum, seminal fluid, amniotic fluid and cerebrospinal fluid. The soluble form of ACE appears to be derived from the membrane-bound form of the enzyme in endothelial cells. A main physiological activity of ACE is that it cleaves the C-terminal dipeptide from angiotensin I to produce the potent vasopressor peptide angiotensin II and inactivates the vasodilatory peptide bradykinin by the sequential removal of two C-terminal dipeptides. As a consequence of the  
30       involvement of ACE in the metabolism of these two vasoactive peptides angiotensin II and bradykinin, ACE has become a crucial molecular target in the treatment of hypertension and congestive heart failure. This has led to the development of highly potent and specific ACE inhibitors which have become clinically important and widespread as orally active drugs to control these conditions of hypertension and congestive heart failure. Whilst the metabolism of vasoactive  
35       peptides remains the best known physiological function of ACE, the enzyme has been also implicated in a range of other physiological processes unrelated to blood pressure regulation such as immunity, reproduction and neuropeptide metabolism due to the localization of ACE and/or the in vitro cleavage of a range of biologically active peptides.

Neutral Endopeptidase (NEP, neprilysin, EC 3.4.24.11) is a zinc metalloprotease and classified as a member of the neprilysin family. NEP was first isolated from the brush border membranes of rabbit kidney. Later, an NEP-like enzyme was identified in rat brain as being involved in the degradation of the opioid peptides, enkephalins. The cloning of the ectoenzyme NEP and subsequent site-directed mutagenesis experiments have shown that, as well as having a similar specificity to thermolysin, it also has a similar active site organization. NEP also shows a thermolysin-like specificity for cleaving peptides on the N-terminal side of hydrophobic residues. With regard to the general distribution of NEP it has been determined in the brain and spinal cord, and lesion and electron microscopic studies generally support a predominantly neuronal localization of NEP, although the enzyme could be present on oligodendrocytes surrounding the fibers of the striato-pallidal and striato-nigral pathways and on Schwann cells in the peripheral nervous system. NEP does not appear to be concentrated on specific membrane interfaces such as the synapse, but is rather uniformly distributed on the surface of neuronal perikarya and dendrites. In the periphery, NEP is particularly abundant in the brush border membranes of the kidney and intestine, the lymph nodes and the placenta, and is found in lower concentrations in many other tissues including the vascular wall of the aorta. By finding that the common acute lymphoblastic leukemia antigen was NEP, it was also shown in the state of the art that the enzyme is transiently present at the surface of lymphohaematopoietic cells and elevated levels are found on mature lymphocytes in certain disease states. The clinical interest in NEP, in particular the interest in NEP inhibitors as potential clinical agents derives from the actions of NEP, in conjunction with another zinc metalloprotease, the aminopeptidase N (APN, membrane alanyl aminopeptidase, EC 3.4.11.2), in degrading the enkephalins and also from its role in degrading atrial natriuretic peptide (ANP). For example, it is known that dual inhibitors of NEP and angiotensin converting enzyme (ACE) are potent antihypertensives, resulting from simultaneously increasing the circulating levels of atrial natriuretic peptide, due to NEP inhibition, and decreasing the circulating levels of angiotensin II, due to ACE inhibition. Further interest in the clinical potential of NEP inhibitors came when the peripheral enzyme was shown to degrade the circulating natriuretic and diuretic peptide, atrial natriuretic peptide. NEP inhibitors were therefore investigated for their antihypertensive properties. From a further example it is known that inhibition of enkephalin metabolism by the synthetic NEP inhibitor, thiorphan, gave naloxone-reversible antinociceptive responses in mice. This opened the possibility that, by increasing the levels of endogenous opioids in the regions of their target receptors, an analgesia could be obtained relatively free of the side-effects of morphine or other classical opiate drugs. It was realized that in order to achieve any significant effect, other enkephalin-metabolizing enzymes also had to be inhibited, in particular the aminopeptidase N (APN). Such dual NEP/APN inhibitors completely block enkephalin metabolism and have strong antinociceptive properties.

Endothelin Converting Enzyme (ECE) catalyses the final step in the biosynthesis of the potent vasoconstrictor peptide endothelin (ET). This involves cleavage of the Trp-Val bond in the inactive intermediate, big-endothelin. ECE-1 is a zinc metalloprotease which is homologous with neutral endopeptidase (NEP; neprilysin; EC 3.4.24.11, see above). Like NEP, ECE-1 is inhibited by the compound phosphoramidon and is a type II integral membrane protein. Unlike NEP, however, ECE-1 exists as a disulfide-linked dimer and is not inhibited by other NEP inhibitors such as thiorphan. Immunocytochemical studies indicate a predominant cell-surface location for ECE-1 where it exists as an ectoenzyme. ECE-1 is localized to endothelial cells and some secretory cells, e.g.  $\beta$ -cells in the pancreas, and in smooth muscle cells. Potent and selective inhibitors of ECE, or dual inhibitors of ECE and NEP, may have therapeutic applications in cardiovascular and renal medicine. Endothelin (ET) which is a 21 amino acid bicyclic peptide containing two intramolecular disulfide bonds, is one of the most potent vasoconstricting peptides identified to date and administration to animals results in a sustained increase in blood pressure emphasizing its potential role in cardiovascular regulation. The endogenous production of ET-1 in humans contributes to the maintenance of basal vascular tone. The endothelin system and related enzymes like ECE therefore represent a likely candidate for the development of novel pharmaceutical agents. Thus, the clinical interest in ECE, in particular the interest in ECE inhibitors as potential clinical agents derives from the actions of ECE, in particular in the context of the biosynthesis of ET. Consequently, compounds showing a significant endothelin converting enzyme inhibitory activity are useful in treating and preventing various diseases which are induced or suspected to be induced by ET.

Particular substrates of metalloproteases or metalloendopeptidases known in the state of the art are e.g. big-endothelin-1 (big-ET-1), atrial natriuretic peptides (ANP), and bradykinin. For example, big-ET-1 is known to be a biologically inactive precursor of endothelin-1 (ET-1) which is a highly potent vasoconstrictor peptide that is produced from its precursor big-endothelin-1 via a specific proteolytic processing. ET-1 has a physiological role in the maintenance of basal vascular tone in humans but also seems to be a causative factor in the pathogenesis of various cardiovascular diseases like hypertension, heart failure and atherosclerosis. One approach to attenuate the adverse effects of ET-1 excess is to inhibit the enzymatic conversion of big-ET-1 to ET-1. Since endothelin converting enzyme-1 (ECE-1) was cloned in 1994 (Xu D. et al., Cell, 1994, 78: 473-485), this enzyme has become generally accepted as the endopeptidase responsible for the physiological conversion of big-ET-1 to ET-1. Also since that time, the NEP-inhibitor phosphoramidon became widely accepted as the tool compound which also potently inhibits ECE-1; furthermore, its activity could be verified in heart failure patients given an infusion of big-ET-1 (Love MP et al, Circ, 1998, 94: 2131-2137).

However, despite the fact that ECE-1 has the ability to cleave big-ET-1, more recent reports raise doubts as to whether ECE-1 is the physiologically relevant endothelin converting enzyme, or at least argue that additional enzymes must be involved in the production of ET-1 (Barker S. et al., Mol Pharmacol, 2001, 59: 163-169). Furthermore, according to Barker et al. endothelin-1 (ET-1) has been implicated as a causative factor in the pathogenesis of hypertension, pulmonary hypertension, congestive heart failure, atherosclerosis, and asthma (see also Douglas, 1997, Trends Pharmacol Sci 18:408-412; Haynes and Web, 1998, J Hypertension 16:1081-1088; Goldie and Henry, 1989, Life Sci 65:1-15). A number of highly potent ET receptor antagonist have been reported for therapeutic use, but these compounds are generally selective for ET<sub>A</sub> receptors or non-selective ET<sub>A</sub>/ET<sub>B</sub> antagonists (Douglas, 1997, supra). Although ET<sub>B</sub> receptors predominate in some tissues, yet they are resistant to blockade by selective ET<sub>B</sub> or non-selective ET<sub>A</sub>/ET<sub>B</sub> antagonists (Hay et al., 1998, J Pharmacol Exp Ther 284:669-677). Therefore, specific inhibition of ET-1 synthesis with ECE inhibitors may be a better approach for attenuating the adverse effects of ET-1 excess under some conditions.

It is an object of the present invention to generate new therapeutic concepts for the treatment and/or prophylaxis of metalloprotease-related diseases, e.g. in a first place by providing therapeutically useful compounds either inhibiting specifically an individual metalloprotease of pharmaceutical interest or specifically inhibiting a selected combination of at least two types of metalloproteases by a of combined mode of action profile; and in a second place by providing therapeutically useful combinations of said metalloprotease inhibiting compounds.

#### SUMMARY OF THE INVENTION

In view of the doubts in the state of the art (Barker S. et al., Mol Pharmacol, 2001, 59: 163-169) as to whether ECE-1 is the only physiologically relevant endothelin converting enzyme, and of the conclusion that additional enzymes must be involved in the production of ET-1, according to the present invention a homology cloning project was performed in order to investigate if so far unknown metalloproteases may play a role in the conversion of big-ET-1 to ET-1, and whether specific inhibitors to the newly identified metalloprotease may inhibit this conversion. These efforts resulted in the discovery of a new human gene with high homology to NEP (54% identity) and a somewhat lower homology to ECE-1 (37% identity) which are the two best characterized members of the neprilysin metalloprotease family. The polypeptide product of this human gene was found to be abundantly expressed in a number of human tissues, and was designated the working title "IGS5" (see co-pending international patent application PCT/EP 00/11532).

Therefore, generally the present invention pertains to the use of a compound having combined, in particular by concurrent, inhibitory activity

- a) on neutral endopeptidase (NEP) and
  - 5 b) on the metalloprotease IGS5 which is a polypeptide comprising an amino acid sequence which has at least 70% identity to one of the amino acid sequences selected from the group of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6;
- or of a pharmaceutically acceptable salt or solvate or biolabile ester thereof, for the manufacture of a medicament (pharmaceutical composition) for treating a mammal, preferably a human,
- 10 suffering from or being susceptible to a disease or condition which can be alleviated or prevented by combined, in particular by concurrent, inhibition of NEP and IGS5.

In a particular aspect the present invention pertains to the use of a compound with combined NEP/IGS5 inhibitory activity, or a pharmaceutically acceptable salt or solvate or biolabile ester thereof, for the manufacture of a medicament (pharmaceutical composition) for

15 treating a mammal, preferably a human, suffering from or being susceptible to a disease or to a condition where big-ET-1 levels are elevated and which disease (condition) can be alleviated or prevented by combined, in particular concurrent, inhibition of NEP and IGS5.

20 In a further particular aspect the present invention pertains to the use of a compound with combined NEP/IGS5 inhibitory activity, or a pharmaceutically acceptable salt or solvate or biolabile ester thereof, for the manufacture of a medicament (pharmaceutical composition) for treating a mammal, preferably a human, suffering from or being susceptible to a disease or condition where ET-1 is significantly upregulated and which disease (condition) can be alleviated

25 or prevented by combined, in particular concurrent, inhibition of NEP and IGS5.

In a further particular aspect the present invention pertains to the use of said compounds with combined or concurrent NEP/IGS5 inhibitory activity or a pharmaceutically acceptable salt or solvate or biolabile ester thereof for the manufacture of a medicament (pharmaceutical composition), preferably for treatment and/or prophylaxis of hypertension, including secondary forms of hypertension such as renal or pulmonary hypertension, heart failure, angina pectoris, arrhythmias, myocardial infarction, cardiac hypertrophy, cerebral ischemia, peripheral vascular disease, subarachnoidal hemorrhage, chronic obstructive pulmonary disease (COPD), asthma, renal disease, atherosclerosis, and pain in colorectal cancer or prostate cancer, in larger

35 mammals, preferably in humans.

Furthermore, it may be beneficial to additionally combine the said compounds showing combined or concurrent NEP/IGS5 inhibitory activity with other individual and/or combined

metalloprotease inhibitors than the NEP/IGS5 inhibitors, e.g. with separate ACE- and/or ECE- and/or NEP-inhibitors and/or mixed inhibitors of these metalloproteases.

## DETAILED DESCRIPTION OF THE INVENTION

5

### Definitions

The terms used in the present application, unless explicitly defined otherwise hereinafter, have the meaning as usually understood by the skilled artisan in this field of invention. In the following some definitions are provided to facilitate understanding of certain terms used frequently herein.

10

"IGS5" refers, among others, to a polypeptide comprising the amino acid sequence set forth in one of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6, or respective variants thereof. Thus "IGS5" particularly includes IGS5PROT, IGS5PROT1 and IGS5PROT2.

15

"Enzyme Activity" or "Biological Activity" refers to the metabolic or physiologic function of said IGS5 including similar activities or improved activities or these activities with decreased undesirable side effects.

20

"IGS5-gene" refers to a polynucleotide comprising the nucleotide sequence set forth in one of SEQ ID NO:1, SEQ ID NO:3 and SEQ ID NO:5, or respective variants, e.g. allelic variants, thereof and/or their complements.

25

"Identity", as known as a measure of identity in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences, e.g. in generally by alignment of the sequences so that the highest order match is obtained. Thus "Identity" and or the alternative wording "Similarity" has an art-recognized meaning and can be readily calculated by known methods, including but not limited to those described in "Computational Molecular Biology", Lesk, A.M., Ed., Oxford University Press, New York, 1988; "Biocomputing: Informatics and Genome Projects", Smith, D.W., Ed., Academic Press, New York, 1993; "Computer Analysis of Sequence Data", Part I, Griffin, A.M., and Griffin, H.G., Eds., Humana Press, New Jersey, 1994; "Sequence Analysis in Molecular Biology", von Heinje, G., Academic Press, 1987; "Sequence Analysis Primer", Gribkov, M. and Devereux, J., Eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., SIAM J. Applied Math., 48: 1073 (1988). Preferred methods to determine identity are designed to give the largest match between the sequences tested. Methods to determine identity

30  
35

and similarity are codified in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387 (1984)), BLASTP, BLASTN, and FASTA (Altschul, S.F. et al., *J. Molec. Biol.* 215: 403-410 (1990)). The BLAST X program is publicly available from NCBI and other sources (BLAST Manual, Altschul, S., et al., NCBI NLM NIH Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.* 215: 403-410 (1990)). The well known Smith Waterman algorithm may also be used to determine identity. A publicly available program useful to determine identity or similarity of polypeptide sequences or polynucleotide sequence, respectively, is known as the "gap" program from Genetics Computer Group, Madison WI, which is usually run with the default parameters for comparisons (along with no penalty for end gaps). The preferred (i.e. default) parameters for polypeptide sequence comparison include the following: Algorithm as described by Needleman and Wunsch, *J. Mol. Biol.* 48: 443-453 (1970); Comparison Matrix BLOSSUM62 from Hentikoff and Hentikoff, *Proc. Natl. Acad. Sci. USA.* 89:10915-10919 (1992); Gap Penalty: 12; Gap Length Penalty: 14. The preferred (i.e. default) parameters for polynucleotide sequence comparison include the following: Algorithm as described by Needleman and Wunsch, *J. Mol Biol.* 48: 443-453 (1970); Comparison Matrix: matches = +10, mismatch = 0; Gap Penalty: 50; Gap Length Penalty: 3. The word "homology" may substitute for the word "identity".

As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 95% "identity" to a reference nucleotide sequence, for example to a reference nucleotide sequence selected from the group of SEQ ID NO:1, SEQ ID NO:3 and SEQ ID NO:5, is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the respective reference nucleotide sequence. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence, or in a number of nucleotides of up to 5% of the total nucleotides in the reference sequence there may be a combination of deletion, insertion and substitution. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

35

Similarly, by a polypeptide having an amino acid sequence having at least, for example 95% "identity" to a reference amino acid sequence, for example to a reference amino acid sequence selected from the group of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6, is intended



that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the respective reference amino acid. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino- or carboxy-terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

"Homolog" is a generic term used in the art to indicate a polynucleotide or polypeptide sequence possessing a high degree of sequence relatedness to a subject sequence. Such relatedness may be quantified by determining the degree of identity and/or similarity between the sequences being compared as herein described. Falling within this generic term are the terms "ortholog", meaning a polynucleotide or polypeptide that is the functional equivalent of a polynucleotide or polypeptide in another species, and "paralog" meaning a functionally similar sequence when considered within the same species. Hence, in humans for example, within the family of endothelin converting enzymes ECE-1 is a paralog of the other members, e.g. of ECE-2.

#### **Pharmacological Evaluation of the Newly identified Metalloprotease IGS5 of Specific Inhibitors to this Metalloprotease**

In view of the doubts in the state of the art (Barker S. et al., Mol Pharmacol, 2001, 59: 163-169) as to whether ECE-1 is the only physiologically relevant endothelin converting enzyme, and of the conclusion that additional enzymes must be involved in the production of ET-1, according to the present invention a homology cloning project was performed in order to investigate if so far unknown metalloproteases may play a role in the conversion of big-ET-1 to ET-1, and whether specific inhibitors to the newly identified metalloprotease may inhibit this conversion. Under the assumption that enzymes with similar activity should have similarities in their amino acid sequences, the human genome was searched for DNA sequences coding for proteins with homology to NEP and ECE, the two best characterized members of the neprilysin metalloprotease family. These efforts resulted in the discovery of a new human gene with high homology to NEP (54% identity) and a somewhat lower homology to ECE-1 (37% identity). The polypeptide product of this gene was found to be abundantly expressed in a number of human tissues, and designated the working title "IGS5". The cloning, expression and basic biological characterization of IGS5 is described in detail in the co-pending international patent application.

PCT/EP 00/11532 which entire content is particularly incorporated by reference to further illustrate the respective details given below on IGS5.

In order to characterize and evaluate the pharmacological enzymatic properties of IGS5  
5 for the purpose of the present invention a human IGS5 protein was generated by using an insect cell line as the expression system, and a variety of potential substrates of the IGS5 protein were tested. IGS5 was confirmed to efficiently cleave big-ET-1, bradykinin and substance P, thus further confirming that this novel protein is a genuine metalloprotease with a broad substrate  
10 specificity, which is a common feature of metalloproteases and which feature has been reported for NEP, ECE-1 and also ACE. It should also be noted that according to the findings of the present invention the proteolysis of big-ET-1 by IGS5 surprisingly results in the correct formation of ET-1, e.g. big-ET-1 is correctly cleaved between amino acids Trp21 and Val22.

Furthermore, according to the present invention for the first time the potency of  
15 metalloprotease inhibitor compounds to suppress the conversion of big-ET to ET-1 was examined, using a labeled fluorescent big-ET-1 analogue. The results are summarized in Table 9 of the experimental section. It is of interest that phosphoramidon that is known to inhibit the conversion of big-ET to ET-1 in vivo, also inhibits IGS5 with high potency in the biochemical assay used in the present invention, and surprisingly that the inhibition of IGS5 by phosphoramidon is actually  
20 considerably higher than ECE-1. In contrast, the selective NEP inhibitor thiorphan as well as the selective ECE-1 inhibitor SM-19712 (4-chloro-N-[[[4-cyano-3-methyl-1-phenyl-1H-pyrazol-5-yl]amino]carbonyl] benzenesulfonamide, monosodium salt; Umekawa K, Hasegawa H, Tsutsumi Y, Sato K, Matsumura Y, Ohashi N., J Pharmacol 2000 Sep;84(1):7-15; Discovery Research Laboratories I, Research Center, Sumitomo Pharmaceuticals Co, Ltd, Osaka, Japan) do not affect  
25 the activity of IGS5 (Table 9, see experimental section).

A very particular aspect of the present invention is the most important and unique finding that numerous compounds which revealed to be metalloprotease inhibitors are able to inhibit IGS5 enzyme even at low nanomolar concentrations, e.g. at concentrations corresponding to IC<sub>50</sub>  
30 values in the range of about 1 to 10 nM, and thus prove to also specifically inhibit the newly identified IGS5 metalloprotease of particular pharmaceutical interest.

Therefore, generally the present invention pertains to the use of a compound having combined, in particular concurrent, inhibitory activity

- 35 a) on neutral endopeptidase (NEP) and  
b) on the metalloprotease IGS5 which is a polypeptide comprising an amino acid sequence which has at least 70% identity to one of the amino acid sequences selected from the group of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6;

or of a pharmaceutically acceptable salt or solvate or biolabile ester thereof, for the manufacture of a medicament (pharmaceutical composition) for treating a mammal, preferably a human, suffering from or being susceptible to a condition which can be alleviated or prevented by combined, in particular concurrent, inhibition of NEP and IGS5.

5

In a particular aspect the present invention pertains to the use of a compound with combined NEP/IGS5 inhibitory activity, or a pharmaceutically acceptable salt or solvate or biolabile ester thereof, for the manufacture of a medicament (pharmaceutical composition) for treating a mammal, preferably a human, suffering from or being susceptible to a disease or condition where big-ET-1 levels are elevated and which disease or condition can be alleviated or prevented by combined, in particular concurrent, inhibition of NEP and IGS5.

10

In a further particular aspect the present invention pertains to the use of a compound with combined NEP/IGS5 inhibitory activity, or a pharmaceutically acceptable salt or solvate or biolabile ester thereof, for the manufacture of a medicament (pharmaceutical composition) for treating a mammal, preferably a human, suffering from or being susceptible to a disease or condition where ET-1 is significantly upregulated and which disease or condition can be alleviated or prevented by combined, in particular concurrent, inhibition of NEP and IGS5.

15

"Combined, or in particular concurrent, inhibitory activity" in the sense of the invention means at least the dual inhibition of NEP and IGS5 by concurrent block of both enzymatic systems, NEP and IGS5, and potentially additional concurrent inhibition of a third system, e.g. triple inhibition of the enzymatic systems NEP, IGS5 and e.g. ECE-1. According to the results of the present invention it may be expected that this combined or concurrent inhibition of both enzymatic systems, NEP and IGS5, is more effective than the isolated blockade of either group by different compounds or just the blockade of each of said individual enzymes. Thus, the present invention provides a new therapeutical concept by suggesting the use of combined, in particular concurrent, NEP and IGS5 inhibitors for the treatment and/or prophylaxis of a set of certain diseases or conditions which can be alleviated or prevented by combined, in particular concurrent, inhibition of NEP and IGS5. In this respect the invention also provides compounds which show combined or concurrent inhibition of both, NEP and IGS5, thereby providing a novel and prospective use of compounds with increased therapeutical value for the treatment and/or prophylaxis of the concerned diseases or conditions.

20

25

30

35

Combined, in particular concurrent, mechanisms of action are of outstanding medical interest as therapeutical benefits can be expected which are more pronounced than modulating each singular system separately.


For example, with regard to endothelin-converting enzyme inhibitors further elucidation of current status and perspectives of this principle and the related benefits recently were reported in a review by B.-M. Löffler (J. Cardiovasc. Pharmacol. (2000), 35(Suppl. 2), S79-S82). According to Löffler, recent research has led to the discovery of potent selective or mixed endothelin-converting enzyme (ECE), ECE/neutral endopeptidase (NEP) and ECE/NEP/angiotensin-converting enzyme (ACE) inhibitors. There is also reported increasing evidence, that the functions of the endothelin (ET), renin-angiotensin and NEP systems for the regulation of the cardiovascular homeostasis are connected by a complex regulation network. Thus Löffler estimates that it will be a challenging task of future research with the newly available selective and mixed ECE-1 inhibitors to show whether the combined inhibition of more than one cardiovascular system is superior to selective inhibition. In this respect the present invention provides a superior progress in that for the first time combined or concurrent inhibition of at least the enzymatic systems NEP and IGS5 is contributed to the state of the art.

In this particular aspect the present invention pertains to the use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof for the manufacture of a medicament (pharmaceutical composition) for treatment and/or prophylaxis of hypertension, including secondary forms of hypertension such as renal or pulmonary hypertension, heart failure, angina pectoris, arrhythmias, myocardial infarction, cardiac hypertrophy, cerebral ischemia, peripheral vascular disease, subarachnoidal hemorrhage, chronic obstructive pulmonary disease (COPD), asthma, renal disease, atherosclerosis, and pain in colorectal cancer or prostate cancer, in mammals, preferably in humans. In particular, in the present invention the compounds with combined or concurrent NEP/IGS5-inhibitory activity preferably are used for the treatment and/or prophylaxis of said diseases or conditions, in a patient sub-population suffering from or being susceptible to a disease or condition which can be alleviated or prevented by combined, in particular concurrent, inhibition of NEP and IGS5.

Furthermore, it may be beneficial to additionally combine the compounds showing combined or concurrent NEP/IGS5 inhibitory activity according to the invention with other individual and/or combined metalloprotease inhibitors than combined NEP/IGS5 inhibitors. Such other metalloprotease inhibitors that may be used in combination with compounds with combined NEP/IGS5 inhibitory activity are for example ACE inhibitors such as captopril, enalapril, lisinopril, fosinopril, perindopril, quinapril, ramipril; furthermore, selective ECE inhibitors such as compound SM-19712 (Sumitomo, supra); selective NEP inhibitors such as thiorphan; dual NEP/ECE inhibitors such as compound CGS-35066 (De Lombart et al., J. Med. Chem. 2000, Feb. 10; 43(3):488-504); or mixed inhibitors of these metalloproteases such as omapatrilat or sampatrilat. By this type of combination treatment and/or prophylaxis the therapeutic value of the compounds with combined or concurrent NEP/IGS5 inhibitory activity still may be further increased, in

particular with regard to the diseases and/or conditions mentioned above. Therefore in a further aspect the invention also pertains to a combination therapy and/or a combination prophylaxis which are further described below.


- 5 In particular, according to the present invention it was found that compounds which are primarily neutral endopeptidase inhibitors (NEP-inhibitors) are well suited to inhibit also the newly identified IGS5 metalloprotease enzyme of pharmaceutical interest even at said low nanomolar concentrations, and thus these compounds prove to be metalloprotease inhibitors with a combined mode of action profile, that is e.g. a combined or concurrent selective NEP/IGS5 inhibitory activity
- 10 profile. Such compounds may have a structure of formula I:




wherein

A stands for a group with formula II or III

15



II



III

in which formula II

R<sup>1a</sup> stands for a phenyl-lower-alkyl group which can be optionally substituted in the phenyl ring by lower alkyl, lower alkoxy or halogen, or for a naphthyl-lower-alkyl group,

- 20 R<sup>2a</sup> means hydrogen or a group forming a biolabile ester; and

in which formula III

R<sup>1b</sup> is hydrogen or a group forming a biolabile phosphonic acid ester,

R<sup>2b</sup> is hydrogen or a group forming a biolabile phosphonic acid ester;

- 25 and wherein

R<sup>3</sup> means hydrogen or a group forming a biolabile carboxylic acid ester; and physiologically acceptable salts of acids or solvates of the formula I.

- 30 Where the substituents in the compounds of formula I are or contain lower alkyl or alkoxy groups, these can be straight-chain or branched and contain, in particular, 1 to 4, preferably 1 to 2,

carbon atoms and are preferably methyl or methoxy. Where the substituents contain halogen, particularly suitable are fluorine, chlorine or bromine, preferably fluorine or chlorine.

5 In the radical  $R^{1a}$  the lower alkylene chain can contain 1 to 4, preferably 1 to 2, carbon atoms.  $R^{1a}$  in particular is an optionally substituted phenethyl group which can optionally be substituted one or more times by halogen, lower alkoxy or lower alkyl, or is a naphthylethyl group.

The compounds of formula Ia are optionally esterified dicarboxylic acid derivatives.

10 Suitable groups  $R^3$  forming biolabile carboxylic acid esters are those which can be cleaved under physiological conditions in vivo with release of the carboxylic acid. For example, those suitable for this purpose are lower alkyl groups, phenyl or phenyl-lower alkyl groups optionally mono- or polysubstituted in the phenyl ring by lower alkyl or lower alkoxy or by a lower alkylene chain bonded to two adjacent carbon atoms, dioxolanylmethyl groups optionally substituted in the dioxolane ring by lower  
15 alkyl or  $C_2$ - $C_8$ -alkanoyloxymethyl groups optionally substituted on the oxymethyl group by lower alkyl. If the group  $R^3$  forming a biolabile ester is or contains lower alkyl, this can be branched or unbranched and can contain 1 to 4 carbon atoms. If the group forming a biolabile ester is an optionally substituted phenyl-lower alkyl group, this can contain an alkylene chain having 1 to 3, preferably 1, carbon atom(s) and is preferably benzyl. If the phenyl ring is substituted by a lower alkylene chain, this can contain 3 to  
20 4, preferably 3, carbon atoms. If  $R^3$  is an optionally substituted alkanoyloxymethyl group, this can contain a preferably branched alkanoyloxy group having 2 to 6, preferably 3 to 5, carbon atoms and can be, for example, a pivaloyloxymethyl radical (= tert-butylcarbonyl-oxymethyl radical).

25 Suitable groups  $R^{2a}$  forming biolabile carboxylic acid esters are those which can be cleaved under physiological conditions in vivo with release of the carboxylic acid, and correspond to the groups exemplified for group  $R^3$  supra.

Groups  $R^{1b}$  and  $R^{2b}$  suitable as groups forming biolabile phosphonic acid esters are those which can be removed under physiological conditions in vivo with release of the respective phosphonic acid function. For example, groups which are suitable for this purpose are lower alkyl groups,  $C_2$ - $C_8$ -alkanoyloxymethyl groups optionally substituted on the oxymethyl group by lower alkyl, or phenyl or phenyl-lower alkyl groups whose phenyl ring is optionally mono- or polysubstituted by lower alkyl, lower  
30 alkoxy or by a lower alkylene chain bonded to two adjacent carbon atoms. If the group  $R^{1b}$  and/or  $R^{2b}$  forming a biolabile ester is or contains lower alkyl, this can be branched or unbranched and can contain 1 to 4 carbon atoms. If  $R^{1b}$  and/or  $R^{2b}$  are an optionally substituted alkanoyloxymethyl group, it can contain a preferably branched alkanoyloxy group having 2 to 6, preferably 3 to 5, carbon atoms and can, for example, be a pivaloyloxymethyl radical (= tert-butylcarbonyloxymethyl radical). If  $R^{1b}$  and/or  
35  $R^{2b}$  are an optionally substituted phenyl-lower alkyl group, this can contain an alkylene chain having 1

to 3, preferably 1, carbon atoms. If the phenyl ring is substituted by a lower alkylene chain, this can contain 3 to 4, in particular 3, carbon atoms and the substituted phenyl ring is in particular indanyl.

5 Suitable physiologically acceptable salts of acids of formula I include their alkali metal, alkaline earth metal or ammonium salts, for example sodium, potassium or calcium salts or salts with physiologically acceptable, pharmacologically neutral organic amines such as, for example, diethylamine or tert-butylamine, or phenyl-lower alkylamines such as  $\alpha$ -methylbenzylamine.


10 The compounds of the formula I contain at least one chiral carbon atom, namely the carbon atom carrying the amide side chain in the 3-position of the benzazepine structure. The compounds can thus be present in two optically active stereoisomeric forms or as a racemate. The present invention includes both the racemic mixtures and the isomerically pure compounds of the formula I. If in the compounds of the formula I the group A stands for formula II, the compounds of formula I contain two chiral carbon atoms, namely the carbon atom which is in position 3 of the  
15 ring framework and carries the amide side-chain, and the carbon atom of the amide side-chain which carries the radical  $R^{1a}$ . These compounds of the formula I, in which the group A stands for formula II, can therefore exist in several optically active stereoisomeric forms or as a racemate. According to the present invention both the racemic mixtures and the isomerically pure compounds may be used. If in the compounds of the formula I the group A stands for formula III,  
20 and  $R^{1b}$  and  $R^{2b}$  are not hydrogen and in each case have different meanings, the phosphorus atom of the phosphonic acid group can also be chiral. The invention also relates to the isomer mixtures and isomerically pure compounds of the formula I, in which the group A stands for formula III, formed as a result of chiral phosphorus atoms.

25 According to the invention, the compounds of the formula I, their salts and biolabile esters may be obtained in a manner known per se in the state of the art (see below).

In preferred embodiment of the present invention it was found that primarily NEP-inhibitory compounds such as benzazepinone-N-acetic acid derivatives with structure of formula  
30 1a, or such as phosphono-substituted benzazepinone derivatives of structure of formula 1b.

Compounds with structure of formula 1a are already known as NEP-inhibiting compounds from US 5,677,297 said compounds being useful for the treatment of diseases or conditions as referenced supra. The compounds of the formula 1a have the following structure  
35

16



1a

wherein

**R<sup>1a</sup>** stands for a phenyl-lower-alkyl group which can be optionally substituted in the phenyl ring by lower alkyl, lower alkoxy or halogen, or for a naphthyl-lower-alkyl group,

5 **R<sup>2a</sup>** means hydrogen or a group forming a biolabile ester and

**R<sup>3</sup>** means hydrogen or a group forming a biolabile ester, and physiologically acceptable salts of acids of the formula I.

10 Where the substituents in the compounds of formula Ia are or contain lower alkyl or alkoxy groups, these can be straight-chain or branched and contain, in particular, 1 to 4, preferably 1 to 2, carbon atoms and are preferably methyl or methoxy. Where the substituents contain halogen, particularly suitable are fluorine, chlorine or bromine, preferably fluorine or chlorine.

15 In the radical **R<sup>1a</sup>** the lower alkylene chain can contain 1 to 4, preferably 1 to 2, carbon atoms. **R<sup>1a</sup>** in particular is an optionally substituted phenethyl group which can optionally be substituted one or more times by halogen, lower alkoxy or lower alkyl, or is a naphthylethyl group.

20 The compounds of formula Ia are optionally esterified dicarboxylic acid derivatives. Depending on the mode of administration, biolabile monoesters, particularly compounds in which **R<sup>2a</sup>** is a group forming a biolabile ester and **R<sup>3</sup>** is hydrogen, or dicarboxylic acids are preferred, the latter being particularly suitable for i.v. administration.

25 Suitable **R<sup>2a</sup>** and **R<sup>3</sup>** groups, in compounds of formula Ia, forming biolabile esters are lower alkyl groups, phenyl or phenyl-lower-alkyl groups which are optionally substituted in the phenyl ring by lower alkyl or by a lower alkylene chain bonded to two adjacent carbon atoms, dioxolanylmethyl groups which are optionally substituted in the dioxolane ring by lower alkyl, or C<sub>2</sub>-C<sub>6</sub>-alkanoyloxymethyl groups optionally substituted on the oxymethyl group by lower alkyl. Where the **R<sup>2a</sup>** or **R<sup>3</sup>** group forming a biolabile ester is lower alkyl, this can be a preferably unbranched alkyl group with 1 to 4, preferably 2, carbon atoms. Where the group forming a biolabile ester is an optionally substituted phenyl-lower-alkyl group, its alkylene chain can contain 1 to 3, preferably 1, carbon atom. Where the phenyl ring is substituted by a lower alkylene chain, this can contain 3 to 4, particularly 3, carbon atoms. Phenyl, benzyl or indanyl are particularly suitable as phenyl-containing substituents **R<sup>2a</sup>** and/or **R<sup>3</sup>**. Where **R<sup>2a</sup>** and/or **R<sup>3</sup>** are an optionally



substituted alkanoyloxymethyl group, their alkanoyloxy group can contain 2 to 6, preferably 3 to 5, carbon atoms and is preferably branched and can be, for example, a pivaloyloxymethyl radical (= tert-butylocarbonyl-oxymethyl radical).

5. Suitable physiologically acceptable salts of dicarboxylic acids or monoesters of formula I include their alkali metal, alkaline earth metal or ammonium salts, for example sodium or calcium salts or salts with physiologically acceptable, pharmacologically neutral organic amines such as, for example, diethylamine or tert-butylamine.

10 The compounds of formula Ia contain two chiral carbon atoms, namely the carbon atom which is in position 3 of the ring framework and carries the amide side-chain, and the carbon atom of the amide side-chain which carries the radical  $R^{1a}$ . The compounds can therefore exist in several optically active stereoisomeric forms or as a racemate. According to the present invention both the racemic mixtures and the isomerically pure compounds of formula Ia may be used.

15

According to the invention, the compounds of the formula Ia and their salts and biolabile esters may be obtained in a manner known per se in the state of the art, e.g. as described in US 5,677,297.

20 Preferred compounds of the formula Ia e.g. those in which  $R^2$  and/or  $R^3$  means a group forming a biolabile ester, and physiologically acceptable salts thereof. The groups forming a biolabile ester may be a lower alkyl group, or a phenyl or phenyl-lower-alkyl group, particularly phenyl, benzyl or indanyl, which is optionally substituted in the phenyl ring by lower alkyl or by a lower alkylene chain bonded to two adjacent carbon atoms, or a dioxolanylmethyl group, particularly (2,2-dimethyl-1,3-dioxolane-4-yl)methyl, which is optionally substituted in the dioxolane  
25 ring by lower alkyl, or a  $C_2$ - $C_6$ -alkanoyloxymethyl group optionally substituted on the oxymethyl group by lower alkyl. In particular compounds of formula Ia are preferred which are characterized in that  $R^2$  is a group forming a biolabile ester and  $R^3$  is hydrogen.

Particular preferred examples of compounds of formula Ia are, e.g.

30 (3S,2'R)-3-{1-[2'-carboxy-4'-phenylbutyl]-cyclopentane-1-carboxylamino}-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetic acid, (compound Ia-1);

(3S,2'R)-3-{1-[2'-(ethoxycarbonyl)-4'-phenylbutyl]-cyclopentane-1-carboxylamino}-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetic acid, (compound Ia-2);

35 (3S,2'R)-3-{1-[2'-(ethoxycarbonyl)-4'-naphthylbutyl]-cyclopentane-1-carboxylamino}-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetic acid, (compound Ia-3);

and physiologically acceptable salts of acids or solvates thereof.

Further particular examples of compounds of formula Ia are, e.g.

3-{1-[2'-(ethoxycarbonyl)-4'-phenylbutyl]-cyclopentane-1-carboxylamino}-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate-tert-butylester, (compound 1a-4);

3-{1-[2'-(ethoxycarbonyl)-4'-phenylbutyl]cyclopentane-1-carboxylamino}-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetic acid, (compound 1a-5);

5 (3S,2'R)-3-{1-[2'-(ethoxycarbonyl)-4'-phenylbutyl]cyclopentane-1-carboxylamino}-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate-tert-butylester, (compound 1a-6);

(3S,2'R)-3-{1-[2'-(carboxy-4'-phenylbutyl]cyclopentane-1-carboxylamino}-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetic acid, (compound 1a-7);

10 3-{1-[2'-(tert-butoxycarbonyl)-4'-phenylbutyl]cyclopentane-1-carboxylamino}-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate-tert-butylester, (compound 1a-8);

3-{1-[2'-(carboxy-4'-phenylbutyl]cyclopentane-1-carboxylamino}-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetic acid, (compound 1a-9);

3-{1-[2'-(tert-butoxycarbonyl)-4'-phenylbutyl]cyclopentane-1-carboxylamino}-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate-benzylester, (compound 1a-10);


15 3-{1-[2'-(carboxy-4'-phenylbutyl]cyclopentane-1-carboxylamino}-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate-benzylester, (compound 1a-11);

3-{1-[2'-(tert-butylcarbonyloxymethoxycarbonyl)-4'-phenylbutyl]cyclopentane-1-carboxylamino}-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate-benzylester, (compound 1a-12);

20 3-{1-[2'-(pivaloyloxymethoxycarbonyl)-4'-phenylbutyl]cyclopentane-1-carboxylamino}-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetic acid, (compound 1a-13);

and physiologically acceptable salts of acids or solvates thereof.

Compounds with structure of formula 1b are already known as NEP-inhibiting compounds and in addition slightly endothelin converting enzyme inhibitors (ECE-inhibitors) from US 5,952,327, said compounds being useful for the treatment of the diseases or conditions referenced supra. Thus, the invention also relates to compounds of the formula 1b



1b

wherein

- 30 R<sup>1b</sup> is hydrogen or a group forming a biolabile phosphonic acid ester,  
 R<sup>2b</sup> is hydrogen or a group forming a biolabile phosphonic acid ester and  
 R<sup>3</sup> is hydrogen or a group forming a biolabile carboxylic acid ester  
 and physiologically acceptable salts of acids of the formula 1b.

The compounds of the formula Ib are acid derivatives comprising carboxylic acid and phosphonic acid groups which are optionally esterified by groups forming biolabile esters. The biolabile esters of the formula Ib are prodrugs of the free acids. Depending on the administration form, the biolabile esters or the acids are preferred, the latter in particular being suitable for i.v. administration.

Groups  $R^{1b}$  and  $R^{2b}$  suitable as groups forming biolabile phosphonic acid esters are those which can be removed under physiological conditions in vivo with release of the respective phosphonic acid function. For example, groups which are suitable for this purpose are lower alkyl groups,  $C_2$ - $C_6$ -alkanoyloxymethyl groups optionally substituted on the oxymethyl group by lower alkyl, or phenyl or phenyl-lower alkyl groups whose phenyl ring is optionally mono- or polysubstituted by lower alkyl, lower alkoxy or by a lower alkylene chain bonded to two adjacent carbon atoms. If the group  $R^{1b}$  and/or  $R^{2b}$  forming a biolabile ester is or contains lower alkyl, this can be branched or unbranched and can contain 1 to 4 carbon atoms. If  $R^{1b}$  and/or  $R^{2b}$  are an optionally substituted alkanoyloxymethyl group, it can contain a preferably branched alkanoyloxy group having 2 to 6, preferably 3 to 5, carbon atoms and can, for example, be a pivaloyloxymethyl radical (= tert-butylcarbonyloxymethyl radical). If  $R^{1b}$  and/or  $R^{2b}$  are an optionally substituted phenyl-lower alkyl group, this can contain an alkylene chain having 1 to 3, preferably 1, carbon atoms. If the phenyl ring is substituted by a lower alkylene chain, this can contain 3 to 4, in particular 3, carbon atoms and the substituted phenyl ring is in particular indanyl.

Suitable groups  $R^3$  for compounds of formula Ib forming biolabile carboxylic acid esters are those which can be cleaved under physiological conditions in vivo with release of the carboxylic acid. For example, those suitable for this purpose are lower alkyl groups, phenyl or phenyl-lower alkyl groups optionally mono- or polysubstituted in the phenyl ring by lower alkyl or lower alkoxy or by a lower alkylene chain bonded to two adjacent carbon atoms, dioxolanylmethyl groups optionally substituted in the dioxolane ring by lower alkyl or  $C_2$ - $C_6$ -alkanoyloxymethyl groups optionally substituted on the oxymethyl group by lower alkyl. If the group  $R^3$  forming a biolabile ester is or contains lower alkyl, this can be branched or unbranched and can contain 1 to 4 carbon atoms. If the group forming a biolabile ester is an optionally substituted phenyl-lower alkyl group, this can contain an alkylene chain having 1 to 3, preferably 1, carbon atom(s) and is preferably benzyl. If the phenyl ring is substituted by a lower alkylene chain, this can contain 3 to 4, preferably 3, carbon atoms. If  $R^3$  is an optionally substituted alkanoyloxymethyl group, this can contain a preferably branched alkanoyloxy group having 2 to 6, preferably 3 to 5, carbon atoms and can be, for example, a pivaloyloxymethyl radical.

According to the invention, the compounds of the formula Ib and their salts and biolabile esters may be obtained in a manner known per se in the state of the art.

Suitable physiologically acceptable salts of acids of the formula Ib are in each case their alkali metal, alkaline earth metal or ammonium salts, for example their sodium, potassium or calcium salts or salts with physiologically acceptable, pharmacologically neutral organic amines such as, for example, diethylamine, tert-butylamine or phenyl-lower alkylamines such as  $\alpha$ -methylbenzylamine.

5

The compounds of the formula Ib contain a chiral carbon atom, namely the carbon atom carrying the amide side chain in the 3-position of the benzazepine structure. The compounds can thus be present in two optically active stereoisomeric forms or as a racemate. The present invention includes both the racemic mixtures and the isomerically pure compounds of the formula I. If R<sup>1b</sup> and R<sup>2b</sup> in compounds of the formula Ib are not hydrogen and in each case have different meanings, the phosphorus atom of the phosphonic acid group can also be chiral. The invention also relates to the isomer mixtures and isomerically pure compounds of the formula I formed as a result of chiral phosphorus atoms.

10

Preferred compound of formula Ib are those, in which R<sup>3</sup> stands for hydrogen or lower alkyl, e.g. C<sub>1</sub>-C<sub>4</sub>-alkyl, in particular C<sub>1</sub>-C<sub>2</sub>-alkyl, and physiologically acceptable salts of acids of the formula Ib.

Particular examples of compounds of formula Ib are, e.g.

20

Benzyl (3S)-3-(1-dibenzylphosphonomethyl-cyclopentane-1-carboxylamino)-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate, (= compound Ib-1);

(3S)-3-(1-Phosphonomethyl-cyclopentane-1-carboxylamino)-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetic acid, (= compound Ib-2);

25

Benzyl (3S)-3-(1-benzylethylphosphonomethyl-cyclopentane-1-carboxylamino)-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate, (= compound Ib-3);

Ethyl (3S)-3-(1-benzylethylphosphonomethyl-cyclopentane-1-carboxylamino)-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate, (= compound Ib-4);

Ethyl (3S)-3-(1-ethylphosphonomethyl-cyclopentane-1-carboxylamino)-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate, (= compound Ib-5);

30

Ethyl (3S)-3-[1-(pivaloyloxymethylethylphosphonomethyl)-cyclopentane-1-carboxylamino]-2,3,4,5-tetrahydro-2-oxo-1H-benzazepine-1-acetate, (= compound Ib-6);

Ethyl (3S)-3-[1-(5-indanylethylphosphonomethyl)-cyclopentane-1-carboxylamino]-2,3,4,5-tetrahydro-2-oxo-1H-benzazepine-1-acetate, (= compound Ib-7);

35

tert-Butyl (3S)-3-(1-benzylethylphosphonomethyl-cyclopentane-1-carboxylamino)-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate, (= compound Ib-8);

Benzyl (3S)-3-(1-ethylphosphonomethyl-cyclopentane-1-carboxylamino)-2,3,4,5-tetrahydro-2-oxo-1H-benzazepine-1-acetate, (= compound Ib-9);

Benzyl (3S)-3-(1-diethylphosphonomethyl-1-cyclopentane-1-carboxylamino)-2,3,4,5-tetrahydro-2-oxo-1H-benzazepine-1-acetate, (= compound Ib-10);

(3S)-3-(1-Diethylphosphonomethyl-cyclopentane-1-carboxylamino)-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetic acid, (= compound Ib-11);

5. Ethyl (3S)-3-(1-diethylphosphonomethyl-cyclopentane-1-carboxylamino)-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate, (= compound Ib-12);

Ethyl (3S)-3-(1-phosphonomethyl-cyclopentane-1-carboxylamino)-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate, (= compound Ib-13);

10. Benzyl (3S)-3-(1-phosphonomethyl-cyclopentane-1-carboxylamino)-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate, (= compound Ib-14);

Benzyl (3S)-3-(1-diisopropylphosphonomethyl-cyclopentane-1-carboxylamino)-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate, (= compound Ib-15);

Ethyl (3S)-3-(1-benzylisopropylphosphonomethyl-cyclopentane-1-carboxylamino)-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate, (= compound Ib-16);

15. tert-Butyl (3S)-3-(1-ethylphosphonomethyl-cyclopentane-1-carboxylamino)-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate, (= compound Ib-17);

tert-Butyl (3S)-3-[1-(pivaloyloxymethyl-ethylphosphonomethyl)-cyclopentane-1-carboxylamino]-2,3,4,5-tetrahydro-2-oxo-1H-benzazepine-1-acetate, (= compound Ib-18);

20. tert-Butyl (3S)-3-(1-phosphonomethyl-cyclopentane-1-carboxylamino)-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-acetate, (= compound Ib-19);

and physiologically acceptable salts of acids thereof.

Particular preferred examples of compounds of formula Ib are, e.g. compound Ib-2, compound Ib-8, compound Ib-18 or compound Ib-19, most preferably compound Ib-8, and  
25. physiologically acceptable salts of acids thereof.

The present invention for the first time provides evidence that, in addition to ECE-1 metalloprotease known previously in the state of the art, the IGS5 type of endothelin converting enzyme also qualifies to be a metalloprotease which is particularly involved in the cleavage of big-ET to ET-1. Therefore, these findings according to the present invention provide new and  
30. interesting prospects regarding improved therapeutical concepts for the treatment and/or prophylaxis of various diseases influenced and/or implied by IGS5 mediated cleavage of big-ET to ET-1, as the present invention suggests the identification and use of therapeutically active compounds that i.a. specifically inhibit IGS5 type metalloprotease, rather than to look for and to  
35. use compounds binding to previously known ECE-1.

It needs to be stressed that the compounds with structure of formula Ia or formula Ib mentioned above were originally selected in the state of the art on the basis of their NEP-inhibitory

activity. Thus, the very high potency towards IGS5 activity in vitro surprisingly found for these compounds according to the present invention is paralleled by the capability of said compounds to substantially lessen the pressor effect of big-ET in anaesthetised rats.

5 In conclusion, the investigations according to the present invention have led to the finding of a novel ECE/NEP-like metalloprotease that efficiently cleaves big-ET and is sensitive not only to the known endothelin converting enzyme inhibitor phosphoramidon, but surprisingly also to a number of defined metalloprotease inhibitors with structures of formula I, preferably with a structure of formula Ia or formula Ib. It is therefore conceivable that IGS5 may play a role in the  
10 production of ET-1, and that compounds with structures such as formula I, preferably such as formula Ia or formula Ib, may exert their in vivo effects by inhibiting this newly identified IGS5 metalloprotease enzyme.

Therefore, further studies elucidating tissue distribution, physiological function and  
15 pathophysiological role of IGS5 in various diseases, preferably in hypertension, renal disease and heart failure, have been initiated in the context of the present invention with compounds showing combined or concurrent NEP/IGS5 inhibitory activity.

In a double-blind placebo-controlled clinical study involving thirteen healthy volunteers  
20 compound Ia-2 dose-dependently inhibited the big-ET-induced pressure response and showed a clear dose related increase in ANP levels, indicative of its NEP-inhibitory properties. ET-1 levels did not increase as would have been expected from a selective NEP inhibitor, whereas big-ET levels were increased dose dependently in the compound Ia-2 groups compared to placebo group, indicating that the breakdown of big-ET was also inhibited. To proof the concept of clinical efficacy  
25 and safety of compound Ia-2 in man, 6 clinical trials were conducted in healthy volunteers and 2 proof of concept trials were conducted in patients: One randomized, placebo-controlled double-blind trial in patients with hypertension (N=191) and one open, baseline controlled pilot trial in patients with congestive heart failure (N=29). Results showed evidence of neutral endopeptidase (NEP) inhibition by significantly increased and sustained plasma concentrations of ANP and it's  
30 second messenger cGMP in both volunteers and patients following oral dosing with compound Ia-2. Furthermore, results in patients with congestive heart failure showed a significant positive correlation between log plasma concentrations of compound Ia-1 (the active metabolite of compound Ia-2) and plasma levels of big-ET following compound Ia-2 administration ( $p < 0.001$ ). In fact plasma levels of big-ET tended to increase following compound Ia-2 administration from  
35 baseline (200mg: 4.9 to 6.5 fmol/ml (+32.6%); 400 mg: 2.3 to 3.5 fmol/ml (+56%)), supporting the concept of activity of orally administered compound Ia-2 to prevent cleavage of big-ET. Most importantly, compound Ia-2 has demonstrated first evidence of significant and clinically relevant anti-hypertensive activity in a recently completed 4-week study (N=191) in patients with WHO

grade I-II hypertension. In the Intent-to-treat patient population (ITT), office diastolic blood pressure versus placebo decreased by -6.9 mmHg (last value under treatment;  $p < 0.001$ ) and office systolic blood pressure versus placebo decreased by -9.2 mmHg (last value under treatment;  $p = 0.003$ ) at the highest dose investigated (200mg bid). This observation is of particular importance as pure NEP inhibitors have been demonstrated to increase ET-1 and consequently rise rather than decrease blood pressure.

Further studies are planned to investigate the dose-response relationship of compound Ia-2 on cardiac hemodynamic parameters in patients with congestive heart failure, and to investigate the time course of anti-hypertensive effects following once daily dosing in patients with hypertension.

#### **IGS5 Metalloproteases in the Context of the Invention**

15. In the context of the present invention reference is made to IGS5 polypeptides (or IGS5 enzymes or IGS5 metalloproteases, e.g. to IGS5PROT, IGS5PROT1 or IGS5PROT2, respectively), in particular to human IGS5 polypeptides (or human IGS5 enzymes). The IGS5 polypeptides may pertain to polypeptides, in particular to human species polypeptides, comprising an amino acid sequence which has at least 70% identity, preferably at least 80% and in particular at least 85 % identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% identity, to one of that selected from the group of SEQ ID NO:2, SEQ ID NO:4 SEQ and SEQ ID NO:6. Such polypeptides include those comprising a IGS5 polypeptide which is identical to one of the amino acid sequences selected from the group of SEQ ID NO:2, SEQ ID NO:4 SEQ and ID NO:6.

25

Such polypeptides also include those IGS5 polypeptides, in particular human IGS5 polypeptides, having an amino acid sequence of at least 70% identity, preferably at least 80% and in particular at least 85 % identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% identity, to one of the amino acid sequences selected from the group of SEQ ID NO:2, SEQ ID NO:4 SEQ and ID NO:6. Such polypeptides include the IGS5 polypeptides which are identical to one of the amino acid sequences selected from the group of SEQ ID NO:2, of SEQ ID NO:4 and SEQ ID NO:6.

Further polypeptides of the present invention include isolated IGS5 polypeptides comprising the sequence contained in one of SEQ ID NO:2, SEQ ID NO:4 SEQ and ID NO:6, and.

The IGS5 polypeptides in the context of the present invention are members of the neprilysin metalloprotease family, and in particular they are human species polypeptides. They are

of interest because several dysfunctions, disorders or diseases have been identified above and in which these newly identified metalloproteases play a critical role in the pathology of the disease.

Thus, according to the present invention it was found that the IGS5 polypeptides may be involved in the metabolism of biologically active peptides, and in particular that these IGS5 polypeptides are metalloprotease type enzymes which may act on a variety of vasoactive peptides. Vasoactive peptides known in the state of the art are e.g. such like atrial natriuretic peptide (ANP), bradykinin, big endothelin (big ET-1), endothelin (ET-1), substance P, and angiotensin-1. Furthermore, it was found that the IGS5 ectodomain, which is a novel human metalloprotease, efficiently hydrolyzes e.g. in vitro a variety of said vasoactive peptides, in particular big-ET-1, bradykinin and substance P.

The IGS5 metalloprotease type enzymes may be inhibited by reference compounds that are used to determine the inhibition properties with regard to enzymes having ECE/NEP-characteristics, e.g. inhibition by compounds such like phosphoramidon. But no inhibition of IGS5 is observed by reference compounds that selectively inhibit NEP, e.g. no inhibition of IGS5 by compounds such as thiorphan, or by reference compounds that selectively inhibit ECE, e.g. no inhibition of IGS5 could be observed for compounds such as SM-19712 (Sumitomo, supra).

Inhibition of IGS5 could be observed at higher concentrations only for reference compounds that inhibit NEP/ECE, e.g. an example is the NEP/ECE inhibitor CGS-35066 (De Lombart et al., supra). The inhibition data of these reference compounds with regard to the inhibition of the IGS5 metalloprotease type enzymes of the present invention are further described in the experimental part below.

The IGS5 polypeptides of the present invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art. Thus, in an example the IGS5 metalloprotease may be generated by methods particularly described in the copending international patent application PCT/EP 00/11532, which is incorporated by reference herein with regard to its entire content, especially with regard to the homology cloning of the human IGS5 gene and to the expression of the corresponding human IGS5 protein.

IGS5 polynucleotides encoding said IGS5 metalloproteases may also be obtained, using standard cloning and screening techniques, from a cDNA library derived from mRNA in cells of human testis tissue, using the expressed sequence tag (EST) analysis (Adam, M.D., et al. Science (1991) 252:1651-1656; Adams, M.D. et al., Nature, (1992) 355:632-634; Adams, M.D., et



al., Nature (1995) 377 Supp:3-174). IGS5 polynucleotides can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques (e.g. F.M. Ausubel et al., 2000, Current Protocols in Molecular Biology).

5           When IGS5 polynucleotides are used for the recombinant production of the IGS5 polypeptides, the polynucleotide may include the coding sequence for the mature polypeptide, by itself; or the coding sequence for the mature polypeptide in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro-  
10           protein sequence, or other fusion peptide portions. For example, a marker sequence which facilitates purification of the fused polypeptide can be encoded. For example, the marker sequence may preferably be a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz et al., Proc Natl Acad Sci USA (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-  
15           translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA. IGS5 polynucleotides which are identical or sufficiently identical to a nucleotide sequence contained in one of SEQ ID NO:1, SEQ ID NO:3 or SEQ ID NO:5, may be used as hybridization probes for cDNA and genomic DNA or as primers for a nucleic acid amplification (PCR) reaction, to isolate full-length cDNAs and genomic clones encoding IGS5 polypeptides and to isolate cDNA and genomic clones of other genes (including genes encoding  
20           paralogs from human sources and orthologs and paralogs from species other than human) that have a high sequence similarity to one of SEQ ID NO:1, SEQ ID NO:3 or SEQ ID NO:5. Typically these nucleotide sequences are at least 70% identical, preferably at least 80% and in particular at least 85 % identical, more preferably at least 90% identical, most preferably at least 95% identical to that of the referent. The probes or primers will generally comprise at least 15 nucleotides,  
25           preferably, at least 30 nucleotides and may have at least 50 nucleotides. Particularly preferred probes will have between 30 and 50 nucleotides. Particularly preferred primers will have between 20 and 25 nucleotides.

          An IGS5 polynucleotide encoding an IGS5 polypeptide, in particular a human IGS5  
30           polypeptide, may be obtained by a process which comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of one of SEQ ID NO: 1, SEQ ID NO:3 or SEQ ID NO:5, or a fragment thereof; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to the skilled artisan. Preferred stringent hybridization conditions  
35           include overnight incubation at 42 °C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate (w/v), and 20 µg/ml denatured, sheared salmon sperm DNA; followed by washing the filters in 0.1 x SSC at about 65 °C. Thus, IGS5 polynucleotides may be obtained by screening

an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of one of SEQ ID NO:1, SEQ ID NO:3 or SEQ ID NO:5, or a fragment thereof.

The skilled artisan will appreciate that, in many cases, an isolated cDNA sequence will be incomplete, in that the region coding for the polypeptide is cut short at the 5' end of the cDNA. This is a consequence of reverse transcriptase, an enzyme with inherently low "processivity" (a measure of the ability of the enzyme to remain attached to the template during the polymerisation reaction), failing to complete a DNA copy of the mRNA template during 1st strand cDNA synthesis. There are several methods available and well known to those skilled in the art to obtain full-length cDNAs, or extend short cDNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (see, for example, Frohman et al., PNAS USA 85, 8998-9002, 1988). Recent modifications of the technique, exemplified by the Marathon<sup>TM</sup> technology (Clontech Laboratories Inc.) for example, have significantly simplified the search for longer cDNAs. In the Marathon<sup>TM</sup> technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an "adaptor" sequence ligated onto each end. Nucleic acid amplification (PCR) is then carried out to amplify the "missing" 5' end of the cDNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using "nested" primers, that is, primers designed to anneal within the amplified product (typically an adaptor specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the known gene sequence). The products of this reaction can then be analyzed by DNA sequencing and a full-length cDNA constructed either by joining the product directly to the existing cDNA to give a complete sequence, or carrying out a separate full-length PCR using the new sequence information for the design of the 5' primer.

Recombinant IGS5 polypeptides may be prepared by processes well known in the art from genetically engineered host cells comprising expression systems which comprise an IGS5 polynucleotide or polynucleotides. Host cells which are genetically engineered with such expression systems may be used for the production of IGS5 polypeptides by recombinant techniques. Cell-free translation systems can also be employed to produce such IGS5 proteins using RNAs derived from IGS5 DNA constructs. Introduction of IGS5 polynucleotides into host cells can be effected by methods described in many standard laboratory manuals, such as Davis et al., Basic Methods in Molecular Biology (1986) and Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989). Such methods include, for instance, calcium phosphate transfection, DEAE-dextran mediated transfection, transfection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection. Representative examples of appropriate hosts include bacterial cells, such as Streptococci, Staphylococci, E. coli, Streptomyces and Bacillus subtilis cells; fungal cells, such as yeast cells and Aspergillus cells;

insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

A great variety of expression systems can be used, for instance, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector which is able to maintain, propagate or express a polynucleotide to produce a polypeptide in a host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook et al., *Molecular Cloning, A Laboratory Manual* (supra). Appropriate secretion signals may be incorporated into the desired polypeptide to allow secretion of the translated protein into the lumen of the endoplasmic reticulum, the periplasmic space or the extracellular environment. These signals may be endogenous to the polypeptide or they may be heterologous signals, i.e. derived from a different species.

20

If an polypeptide is to be expressed for use in screening assays, Generally it is possible that the IGS5 polypeptide is produced at the surface of the cell or alternatively in a soluble protein form. If the IGS5 polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the IGS5 polypeptide. If produced intracellularly, the cells must first be lysed before the IGS5 polypeptide is recovered. If the IGS5 polypeptide is bound at the surface of the cell (membrane bound polypeptide), usually membrane fractions are prepared in order to accumulate the membrane bound IGS5 polypeptide. IGS5 polypeptides can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during intracellular synthesis, isolation and or purification.

35

Isolated IGS5 polynucleotides, in particular isolated human IGS5 polynucleotides, that may be used to generate an IGS5 polypeptide usually comprise a nucleotide sequence that has at least 70% identity, preferably at least 80% and in particular at least 85 % identity, more preferably

at least 90% identity, yet more preferably at least 95% identity, to a nucleotide sequence encoding one of the polypeptides selected from the group of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6, over the entire coding region. In this regard, polynucleotides which have at least 97% identity are highly preferred, whilst those with at least 98-99% identity are more highly preferred, and those with at least 99%, in particular 99.9%, identity are most highly preferred. For example, such isolated, in particular human, IGS5 polynucleotides that may be used to generate IGS5 polypeptides include nucleotide sequences which have at least 70% identity, preferably at least 80% and in particular at least 85 % identity, more preferably at least 90% identity, yet more preferably at least 95% identity, over the entire length to one of the nucleotide sequences selected from the group of SEQ ID NO:1, SEQ ID NO:3 and SEQ ID NO: 5. In this regard, IGS5 polynucleotides which comprise or have a nucleotide sequence of at least 97% identity to one of the nucleotide sequences selected from the group of SEQ ID NO:1, SEQ ID NO:3 and SEQ ID NO:5 are highly preferred, whilst those with at least 98-99% identity, are more highly preferred, and those with at least 99%, in particular 99.9%, identity are most highly preferred. The IGS5 polynucleotide sequence of SEQ ID NO:1 (designated "IGS5DNA") is indicated in Table 1 representing a cDNA sequence from human origin (Homo sapiens) with a length of 2076 nucleotides and comprises a polypeptide encoding sequence (from nucleotide no. 1 to no. 2073) encoding a polypeptide of 691 amino acids, the polypeptide of SEQ ID NO:2 (designated "IGS5PROT") which is indicated in Table 2. The nucleotide sequence of SEQ ID NO:3 (designated "IGS5DNA1") is indicated in Table 3 representing a cDNA sequence from human origin (Homo sapiens) with a length of 2340 nucleotides (including the stop codon tag) and comprises a polypeptide encoding sequence (from nucleotide no. 1 to no. 2337) encoding a polypeptide of 779 amino acids, the polypeptide of SEQ ID NO:4 (designated "IGS5PROT1") which is indicated in Table 4. The nucleotide sequence of SEQ ID NO:5 (designated "IGS5DNA2") is indicated in Table 5 representing a cDNA sequence from human origin (Homo sapiens) with a length of 2262 nucleotides (including the stop codon tag) and comprises a polypeptide encoding sequence (from nucleotide no. 1 to no. 2259) encoding a polypeptide of 753 amino acids, the polypeptide of SEQ ID NO:6 (designated "IGS5PROT2") which is indicated in Table 6.

### **Compounds with Combined or Concurrent Selective NEP/IGS5-Inhibitory Activity as New Therapeutical Concept**

The findings of the present invention have shown that IGS5 metalloproteases, e.g. also in combination with at least one other metalloprotease such as in particular NEP, and optionally in addition ECE and/or ACE, are responsible for one or more biological functions related to the diseases mentioned herein before. Thus, in its broadest aspect the invention generally provides new therapeutic concepts for the treatment of said diseases, as stated already above, by suggesting for the first time to use compounds with combined or concurrent inhibitory activity on

neutral endopeptidase (NEP) and on the metalloprotease IGS5, or a pharmaceutically acceptable salt or solvate or biolabile ester thereof, for the manufacture of a medicament (pharmaceutical composition) for treating a larger mammal, preferably a human, suffering from or being susceptible to a condition which can be alleviated or prevented by combined or concurrent inhibition of NEP and IGS5.

Such compounds useful according to the invention in that they concurrently inhibit the function of the IGS5 metalloprotease and of NEP may be identified by screening methods using IGS5 metalloprotease, and optionally NEP, in an appropriate enzyme inhibition assay format. Such enzyme inhibition assay formats are described in more detail in the experimental section below. For identification of compounds with combined or concurrent selective NEP/IGS5-inhibitory activity candidate compounds may be tested separately in both, an NEP-inhibition assay and IGS5-inhibition assay. NEP/IGS5-inhibitory compounds may be identified from a variety of sources, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures.

The screening method may simply measure the influence of a candidate compound on the activity of the polypeptide excreted into a culture medium, or on cells or membranes bearing the polypeptide. Alternatively, the screening method may involve competition with a competitor. Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the activity of the polypeptide excreted into a culture medium or to the cells or membranes bearing the polypeptide. Inhibition of polypeptide activity is generally assayed in the presence of a known substrate and the effect of the candidate compound is observed by altered activity, e.g. by testing whether the candidate compound results in inhibition of the polypeptide. For example, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide of interest in the context of the present invention, and a suitable substrate to form a mixture, measuring the polypeptide activity in the mixture, and comparing the polypeptide activity of the mixture to a standard without candidate compound.

The present invention also enables the person skilled in the art to identify compounds, e.g. candidate compounds, by means of screening methods involving the findings of the present invention, said compounds may reveal as prospective drug candidates in particular with respect to dysfunctions, disorders or diseases that are referenced already above. It will be readily appreciated by the skilled artisan that an IGS5 metalloprotease may also be used in a method for the structure-based design of IGS5 inhibitory compounds, by:

- (a) determining or using in the first instance the three-dimensional structure of the IGS5 metalloprotease;

(b) deducing the three-dimensional structure for the likely reactive or binding site(s) of an IGS5 inhibitor;

(c) synthesizing candidate compounds that are predicted to bind to or react with the deduced binding or reactive site; and

5 (d) testing whether the candidate compounds are indeed IGS5 inhibitors.

It will be further appreciated that this will normally be an iterative process.

Today, medicinal chemists are well aware of modern strategies for planning and performing organic synthesis in order to generate new substances or compounds that are worth to be  
10 investigated for potential physiological or pharmacological properties, and which compounds therefore promise to prove as prospective new drug candidates for the treatment and/or prophylaxis of specific dysfunctions, disorders or diseases. Furthermore, today it is common to provide compound libraries by means of combinatorial chemistry, e.g. in particular of general and  
15 of "directed" chemical or compound libraries, in which the structure and the variations of pharmacophore groups and the residues or substituents are known to the concerned artisan. If chemical libraries or compound libraries with still unknown structure of the compounds are investigated in screening assays, potential prospective compounds, e.g. candidate compounds, nevertheless, may easily be analysed in their structure and chemical properties by today's well-  
20 established analytical means such as e.g. mass spectroscopy, nuclear magnetic resonance, infrared spectra, melting points, optical rotation if chiral compounds are involved, and elemental analysis.

Thus the invention also pertains to a process for preparing a candidate compound with a defined chemical structure capable of inhibiting the IGS5 polypeptide, said process is comprising  
25 the manufacture of a compound or of a pharmaceutically acceptable salt or biolabile ester thereof by means of chemical synthesis, provided that the activity of the compound to inhibit the IGS5 polypeptide is identifiable by a screening method, e.g. such as described in the experimental section of the present invention.

30 For details of e.g. chemical organic synthesis, and e.g. chemical, analytical and physical methods see the Handbook "Houben-Weyl" (Houben-Weyl, "Methoden der organischen Chemie", Georg Thieme Verlag, Stuttgart, New York) in its most recent version.

One embodiment of the present invention pertains to the use of a compound of formula I  
35 as given supra or a pharmaceutically acceptable salt or solvate or biolabile ester thereof, for the manufacture of a medicament (pharmaceutical composition) for treating a larger mammal, preferably a human, suffering from or being susceptible to a condition which can be improved or prevented by combined or concurrent inhibition of

- a) neutral endopeptidase (NEP) and
- b) of the metalloprotease IGS5 which is a polypeptide comprising an amino acid sequence which has at least 70% identity over the entire length to one of the amino acid sequences selected from the group of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO: 6.

5

A preferred embodiment of the present invention pertains to the use of a compounds according to the invention, in particular compounds with formula I, having combined or concurrent inhibitory activity on

- a) neutral endopeptidase (NEP) and
  - 10 b) on the metalloprotease IGS5 which is a polypeptide comprising an amino acid sequence which has at least 70% identity over the entire length to one of the amino acid sequences selected from the group of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6;
- or a pharmaceutically acceptable salt or solvate or biolabile ester thereof, for the manufacture of a medicament (pharmaceutical composition) for treatment and/or prophylaxis of hypertension,
- 15 including secondary forms of hypertension such as renal or pulmonary hypertension, heart failure, angina pectoris, arrhythmias, myocardial infarction, cardiac hypertrophy, cerebral ischemia, peripheral vascular disease, subarachnoidal hemorrhage, chronic obstructive pulmonary disease (COPD), asthma, renal disease, atherosclerosis, and pain in colorectal cancer or prostate cancer, in larger mammals, preferably in humans.

20

Compounds of formula I can be prepared according to the disclosure in US 5,677,297 which is suitable for compounds of formula Ia and according to US 5,952,327 which is suitable for compounds of formula Ib.

## 25 Protein-Ligand Complexes in Drug Design and Lead Structure optimization

- In another aspect the invention relates to a protein-ligand-complex comprising an IGS5 polypeptide of at least 70% identity to one of the polypeptides of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:5 and an IGS5-binding compound, preferably a compound with IGS5-inhibitory activity of at least that of or being comparable to that of compounds of formula I. Such protein-
- 30 ligand-complexes are particularly useful in drug design methods, lead structure finding, lead structure optimization and modulation methods. The methods are well known in the state of the art. For exemplary reference see literature concerning e.g. combinatorial synthesis and multidimensional NMR-spectroscopy and its contribution to the understanding of protein-ligand-interactions (Kessler, Angew. Chem. 1997, 109, 857-859; James K. Chen et al., Angew. Chem. 107 (1995), S. 1041-1058). Furthermore see Fesik (Journal of Medicinal Chemistry, 34 (1991), S. 2937-2945) who describes NMR studies of molecular complexes as a tool in drug design; and .
- 35 Fesik et al. (Bi chemical Pharmacology 40 (1990), S. 161-167) who describe NMR methods for determining the structures of enzyme/inhibitor complexes as an aid in drug design. A very recent

report of Ross et al. (Journal of Biomolecular NMR, 16: 139-146 (2000)) describes the automation of NMR measurements and data evaluation for systematically screening interactions of small molecules with target proteins, e.g. receptors.

5 Thus, the invention also pertains to the use of a protein-ligand-complex comprising an IGS5 polypeptide of at least 70% identity to one of the polypeptides of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6 and an IGS5-binding compound for the design and modulation or optimization of lead structures with IGS5-binding and IGS5-inhibitory activity.

#### 10 **Combination Therapy (NEP/IGS5-Inhibitory Compounds and ECE / ACE-Inhibitory Compounds)**

As already shortly addressed supra, it may be beneficial to additionally combine compounds showing combined or concurrent NEP/IGS5 inhibitory activity according to the invention, e.g. compounds of formula I, preferably compounds of formula Ia or Ib, with other  
15 individual and/or combined metalloprotease inhibitors than combined NEP/IGS5 inhibitors. Such other metalloprotease inhibitors that may be used in combination with said compounds with combined NEP/IGS5 inhibitory activity are for example ACE inhibitors such as captopril, enalapril, lisinopril, fosinopril, perindopril, quinapril, ramipril; furthermore, selective ECE inhibitors such as compound SM-19712 (Sumitomo, supra); selective NEP inhibitors such as thiorphan; dual  
20 NEP/ECE inhibitors such as compound CGS-35066 (De Lombart et al., J. Med. Chem. 2000, Feb. 10; 43(3):488-504); or mixed inhibitors of these metalloproteases such as omapatrilat or sampatrilat. By this type of combination treatment and/or prophylaxis the therapeutic value of the compounds with combined or concurrent NEP/IGS5 inhibitory activity still may be further increased, in particular with regard to the diseases and/or conditions mentioned above. Therefore  
25 in a further aspect the invention particularly also pertains to a combination therapy and/or combination prophylaxis. By this type of combination treatment and/or prophylaxis the therapeutic value of said compounds with combined or concurrent NEP/IGS5 inhibitory activity, in particular of compounds with formula I, preferably of compounds with formula Ia or Ib, still may be further increased, in particular with regard to the diseases and/or conditions mentioned above.

30

Thus, in this respect the invention pertains to the use of a first compound showing combined or concurrent NEP/IGS5 inhibitory activity or a pharmaceutically acceptable salt or solvate or biolabile ester thereof, as these are described above with regard to the present invention, in combination with at least one additional compound selected from the group of other  
35 individual and/or combined metalloprotease inhibitors than the combined NEP/IGS5 inhibitors, said additional compound preferably being selected from the group of ACE inhibitors, selective ECE inhibitors, selective NEP inhibitors, dual NEP/ECE inhibitors, and mixed inhibitors of these metalloproteases, for the manufacture of a medicament (pharmaceutical composition) for



combination treatment and/or combination prophylaxis of any of the diseases or conditions as referenced above in the context of the present invention. Particularly this use according to the present invention of said first compound in combination with at least one of said additional compounds, is characterized in that the first compound has a structure of formula I, preferably a structure of formula Ia or of formula Ib, as these formulas are referenced above in the context of the present invention. Preferably, the use of said first compound in combination with at least one of said additional compounds, is further characterized in that the combination is co-effective, preferably synergistically effective.

Furthermore, the invention in this respect pertains to pharmaceutical composition (medicament), comprising co-effective, preferably synergistically effective, amounts of:

a first compound with combined or concurrent NEP/IGS5 inhibitory activity or a pharmaceutically acceptable salt or solvate or biolabile ester thereof, as these are described above with regard to the present invention; and of at least one additional compound selected from the group of other individual and/or combined metalloprotease inhibitors than the combined NEP/IGS5 inhibitors, said additional compound preferably being selected from the group of ACE inhibitors, selective ECE inhibitors, selective NEP inhibitors, dual NEP/ECE inhibitors, and mixed inhibitors of these metalloproteases, for combination treatment and/or combination prophylaxis of any of the diseases or conditions as referenced above in the context of the present invention. In particular the pharmaceutical composition according to the present invention may comprise co-effective, preferably synergistically effective, amounts of said first compound and of at least one of said additional compounds, being further characterized in that the first compound has a structure of formula I, preferably a structure of formula Ia or of formula Ib, as these formulas are referenced above in the context of the present invention.

It is self-explaining to the skilled person that combination therapy and/or combination prophylaxis according to the present invention may be achieved by administering to a patient in need of such a therapy and/or such prophylaxis the first compound with combined or concurrent NEP/IGS5 inhibitory activity or a pharmaceutically acceptable salt or solvate or biolabile ester thereof and the additional compound selected from the group of other individual and/or combined metalloprotease inhibitors than the combined NEP/IGS5 inhibitors, in a simultaneous manner, either by administering a single pharmaceutical combination preparation or by separate pharmaceutical preparation for the first and the second compound, in a separate manner, e.g. under a given dosage regimen or scheme which may be either continuous or sequential, or in a graded manner, whatever seems suitable with regard to the patient's disease or condition to be alleviated and/or prevented.

### Formulation and Administration

Th foregoing findings according to the invention show that combined or concurrent selective NEP/IGS5-inhibitory compounds, optionally in combination with separat ACE- and/or ECE-inhibitory compounds, or their respective pharmaceutically acceptable salts or solvates or biolabile esters thereof exert a beneficial therapeutic activity. Therefore these compounds, optionally in compination with separate ACE- and/or ECE-Inhiibitors, are suitable as medicaments for the treatment and/or prophylaxis of hypertension, including secondary forms of hypertension such as renal or pulmonary hypertension, heart failure, angina pectoris, arrhythmias, myocardial infarction, cardiac hypertrophy, cerebral ischemia, peripheral vascular disease, subarachnoidal hemorrhage, chronic obstructive pulmonary disease (COPD), asthma, renal disease, atherosclerosis, and pain in colorectal cancer or prostate cancer, in larger mammals, especially in humans. NEP/IGS5-inhibitory compounds, optionally in combination with separate ACE- and/or ECE-inhibitory compounds, may be given by all known administration routes.

The composition will be adapted to the route of administration, for instance by a systemic or an oral route. Preferred forms of systemic administration include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition compounds can be formulated in an enteric or an encapsulated formulation, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels, and the like.

For administration according to the invention the therapeutically active quantities of the NEP/IGS5-inhibitory compounds that alleviate and/or prevent the diseases or conditions mentioned supra in the context of the invention can be contained together with customary pharmaceutical excipients and/or additives in solid or liquid pharmaceutical formulations.

Examples of solid dosage forms are such as solid, semi-solid, lyophilized powder, tablets, coated tablets, pills, capsules, powders, granules or suppositories, also in form of sustained release formulations. These solid dosage forms can contain standard pharmaceutical inorganic and/or organic excipients. Such excipients include pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, gelatine, sucrose, magnesium carbonate, and the like in addition to customary pharmaceutical additives such as fillers, lubricants or tablet disintegrants. Liquid preparations such as solutions, suspensions or emulsions of the active ingredients can contain the usual diluents such as water, oil and/or suspending aids such as poly thylene glycols and such like. Further additives such as preservatives, flavouring agents and such like may also be added.

The active ingredients can be mixed and formulated with the pharmaceutical excipients and/or additives in a known manner. For the manufacture of solid dosage forms, for example, the active ingredients may be mixed with the excipients and/or additives and granulated in a wet or dry process. Granules or powder can be filled directly into capsules or compressed into tablet cores. If desired, these can be coated in the known manner.

Liquid preparations can be prepared by dissolving or dispersing the compounds and optional pharmaceutical adjuvants, in a carrier, such as, for example, aqueous saline, aqueous dextrose, glycerol, or ethanol, to form a solution or suspension.

The doses to be administered may differ between individuals and naturally vary depending on the type of condition to be treated and the route of administration. For example, locally applicable formulations injectable formulations, generally contain substantially less amount of active substance than systemically applicable formulations. Thus, the dosage range required depends on the judgment of the attending practitioner, in particular in view of the choice of compounds, the route of administration, the nature of the formulation, and the nature of the subject's condition. Suitable dosages, however, are in the range of 0.1-100 µg/kg of subject. Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

25

#### Tables with IGS5 DNA and IGS5 protein sequences

Table 1: IGS5-DNA ("IGS5DNA") of SEQ ID NO:1

5' -  
TGCACCACCCCTGGCTGCGTGATAGCAGCTGCCAGGATCCTCCAGAACATGGACCCGACC  
ACGGAACCGTGTGACGACTTCTACCAGTTTGCATGCGGAGGCTGGCTGCGGCGCCACGTG  
ATCCCTGAGACCAACTCAAGATACAGCATCTTTGACGTCCTCCGCGACGAGCTGGAGGTC  
ATCCTCAAAGCGGTGCTGGAGAATTCGACTGCCAAGGACCGCGGCTGTGGAGAAGGCC  
AGGACGCTGTACCGCTCCTGCATGAACCAGAGTGTGATAGAGAAGCGAGGCTCTCAGCCC  
CTGCTGGACATCTTGGAGGTGGTGGGAGGCTGGCCGGTGGCGATGGACAGGTGGAACGAG  
ACCGTAGGACTCGAGTGGGAGCTGGAGCGGCAGCTGGCGCTGATGAACTCACAGTTCAAC

AGGCGCGTCCTCATCGACCTCTTCATCTGGAACGACGACCAGAACTCCAGCCGGCACATC  
ATCTACATAGACCAGCCACCTTGGGCATGCCCTCCCGAGAGTACTACTTCAACGGCGGC  
AGCAACCGGAAGGTGCGGGAAGCCTACCTGCAGTTCATGGTGTCACTGGCCACGTTGCTG  
CGGGAGGATGCAAACCTGCCCAGGGACAGCTGCCTGGTGCAGGAGGACATGATGCAGGTG  
CTGGAGCTGGAGACACAGCTGGCCAAGGCCACGGTACCCCAGGAGGAGAGACACGACGTC  
ATCGCCTTGTACCACCGGATGGGACTGGAGGAGCTGCAAAGCCAGTTTGGCCTGAAGGGA  
TTTAACTGGACTCTGTTTATACAAACTGTGCTATCCTCTGTCAAATCAAGCTGCTGCCA  
GATGAGGAAGTGGTGGTCTATGGCATCCCCTACCTGCAGAACCTTGAAAACATCATCGAC  
ACCTACTCAGCCAGGACCATAACAGAACTACCTGGTCTGGCGCCTGGTGTGCTGGACCGCATT  
GGTAGCCTAAGCCAGAGATTCAAGGACACACGAGTGAACCTACCGCAAGGCGCTGTTTGGC  
ACAATGGTGGAGGAGGTGCGCTGGCGTGAATGTGTGGGCTACGTCAACAGCAACATGGAG  
AACGCCGTGGGCTCCCTCTACGTCAGGGAGGCGTTCCCTGGAGACAGCAAGAGCATGGTC  
AGAGAACTCATTGACAAGGTGCGGACAGTGTTTGTGGAGACGCTGGACGAGCTGGGCTGG  
ATGGACGAGGAGTCCAAGAAGAAGGCGCAGGAGAAGGCCATGAGCATCCGGGAGCAGATC  
GGGCACCCTGACTACATCCTGGAGGAGATGAACAGGCGCCTGGACGAGGAGTACTCCAAT  
CTGAACTTCTCAGAGGACCTGTACTTTGAGAACAGTCTGCAGAACCTCAAGGTGGGCGCC  
CAGCGGAGCCTCAGGAAGCTTCGGGAAAAGGTGGACCCAAATCTCTGGATCATCGGGGCG  
GCGGTGGTCAATGCGTTCTACTCCCCAAACCGAAACCAGATTGTATTCCCTGCCGGGATC  
CTCCAGCCCCCTTCTTCAGCAAGGAGCAGCCACAGGCCTTGAACTTTGGAGGCATTGGG  
ATGGTGATCGGGCAGGATCACGCACGGCTTTGACGACAATGGCCGGAACCTTCGACAAG  
AATGGCAACATGATGGATTGGTGGAGTAACTTCTCCACCCAGCACTTCCGGGAGCAGTCA  
GAGTGCATGATCTACCAGTACGGCAACTACTCCTGGGACCTGGCAGACGAACAGAACGTG  
AACGGATTCAACACCCTTGGGGAAAACATTGCTGACAACGGAGGGGTGCGGCAAGCCTAT  
AAGGCCTACCTCAAGTGGATGGCAGAGGGTGGCAAGGACCAGCAGCTGCCCGGCCTGGAT  
CTCACCCTATGAGCAGCTCTTCTTCATCAACTACGCCCAGGTGTGGTGGGGTCTTACCGG  
CCCGAGTTCGCCATCCAATCCATCAAGACAGACGTCCACAGTCCCCTGAAGTACAGGGTA  
CTGGGGTGCCTGCAGAACCTGGCCGCCTTCGCAGACACGTTCCACTGTGCCCGGGGCACC  
CCCATGCACCCCAAGGAGCGATGCCGCGTGTGGTAG - 3'

Table 2: IGS5-protein ("IGS5PROT") of SEQ ID NO:2

CTTPGCVIAAARILQNMPTTEPCDDFYQFACGGWLRHVIPETNSRYSIFDVLRLDELEV  
ILKAVLENSTAKDRPAVEKARTLYRSCMNQSVIEKRGSQPLLDILEVVGGWPVAMDRWNE  
TVGLEWELERQLALMNSQFNRRVLIDLFIWNDDQNSSRHIIYIDQPTLGMPSREYYFNQG  
SNRKVREAYLQFMVSVATLLREDANLPRDSCLVQEDMMQVLELETQLAKATVPQERHDV  
IALYHRMGLEELQSQFGLKGFNWTLFITQTVLSSVKIKLLPDEEVVYGIPLYLQNLNIID  
TYSARTIQNYLVWRLVLDRIQSLQRFKDRVNYRKALFGTMVEEVRWRECVGYVNSNME  
NAVGSlyvREAFPGDSKSMVRELIDKVRTVFVETLDELGWMDEESKKKAQEKAMSIREQI

GHPDYILEEMNRRLDEEYSNLNFSEDLYFENSLQNLKVGAQRSRLKLRKVDPNLWIIGA  
AVVNAFYSPNRNQIVFPAGILQPPFFSKEQPQALNFGGIGMVGHEITHGFDDNGRNF DK  
NGNMMDWWSNFSTQHFREQSECMYQYGNYSWDLADEQNVNGFNTLGENIADNGGVRQAY  
KAYLKWMAEGGKDQQLPGLDLTHEQLFFINYAQVWCGSYRPEFAIQSIKTDVHSPKLYRV  
LGSLQNLAAFADTFHCARGTPMHPKERCVRW

Table 3: IGS5-DNA-1 ("IGS5DNA1") of SEQ ID NO: 3

5' -

ATGGGGAAGTCCGAAGGCCCGTGGGGATGGTGGAGAGCGCTGGCCGTGCAGGGCAGAAG  
CGCCCGGGGTTCTCTGGAGGGGGGGCTGCTGCTGCTGCTGCTGCTGGTGACCGCTGCCCTG  
GTGGCCTTGGGTGTCCTCTACGCCGACCGCAGAGGGAAGCAGCTGCCACGCCTTGCTAGC  
CGGCTGTGCTTCTTACAGGAGGAGAGGACCTTTGTAAAACGAAAACCCCGAGGGATCCCA  
GAGGCCCAAGAGGTGAGCGAGGTCTGCACCACCCCTGGCTGCGTGATAGCAGCTGCCAGG  
ATCCTCCAGAACATGGACCCGACCACGGAACCGTGTGACGACTTCTACCAGTTTGCATGC  
GGAGGCTGGCTGCGGCGCCACGTGATCCCTGAGACCAACTCAAGATACAGCATCTTTGAC  
GTCCTCCGCGACGAGCTGGAGGTTCATCCTCAAAGCGGTGCTGGAGAATTGCACTGCCAAG  
GACCGGCCGGCTGTGGAGAAGGCCAGGACGCTGTACCGCTCCTGCATGAACCAGAGTGTG  
ATAGAGAAGCGAGGCTCTCAGCCCCCTGCTGGACATCTTGGAGGTGGTGGGAGGCTGGCCG  
GTGGCGATGGACAGGTGGAACGAGACCGTAGGACTCGAGTGGGAGCTGGAGCGGCAGCTG  
GCGCTGATGAACTCACAGTTCAACAGGCGCGTCCTCATCGACCTCTTCATCTGGAACGAC  
GACCAGAACTCCAGCCGGCACATCATCTACATAGACCAGCCACCTTGGGCATGCCCTCC  
CGAGAGTACTACTTCAACGGCGGCAGCAACCGGAAGGTGCGGGAAGCCTACCTGCAGTTC  
ATGGTGTCAAGTGGCCACGTTGCTGCGGGAGGATGCAAACCTGCCCAGGGACAGCTGCCTG  
GTGCAGGAGGACATGATGCAGGTGCTGGAGCTGGAGACACAGCTGGCCAAGGCCACGGTA  
CCCCAGGAGGAGAGACACGACGTCATCGCCTTGTAACCGGATGGGACTGGAGGAGCTG  
CAAAGCCAGTTTGGCCTGAAGGGATTAACTGGACTCTGTTTATACAAACTGTGCTATCC  
TCTGTCAAAATCAAGCTGCTGCCAGATGAGGAAGTGGTGGTCTATGGCATCCCCTACCTG  
CAGAACCTTGAAAACATCATCGACACCTACTCAGCCAGGACCATAACAGAACTACCTGGTC  
TGGCGCCTGGTGCTGGACCGCATTTGGTAGCCTAAGCCAGAGATTCAGGACACACGAGTG  
AACTACCGCAAGGCGCTGTTTGGCACAATGGTGGAGGAGGTGCGCTGGCGTGAATGTGTG  
GGCTACGTCAACAGCAACATGGAGAACGCCGTGGGCTCCCTCTACGTGAGGAGGCGTTC  
CCTGGAGACAGCAAGAGCATGGTCAGAGAACTCATTGACAAGGTGCGGACAGTGTTTGTG  
GAGACGCTGGACGAGCTGGGCTGGATGGACGAGGAGTCCAAGAAGAAGGCGCAGGAGAAG  
GCCATGAGCATCCGGGAGCAGATCGGGCACCCCTGACTACATCCTGGAGGAGATGAACAGG  
CGCCTGGACGAGGAGTACTCCAATCTGAACTTCTCAGAGGACCTGTACTTTGAGAACAGT  
CTGCAGAACCTCAAGGTGGGCGCCCAGCGGAGCCTCAGGAAGCTTCGGGAAAAGGTGGAC  
CCAAATCTCTGGATCATCGGGGCGGCGGTGGTCAATGCGTTCCTACTCCCCAAACCGAAAC

- 3 -

MKGSEGPVGMVESAGRAGQKRPGFLEGGLLLLLLLLLVTAALVALGVLYADRRGKQLPRLAS  
RLCFLQEERTFVVKRKPGRGPEAQEVSEVCTTPGCVIAAARILQNMDPTEPCDDFYQFAC  
GGWLRHHVIPETNSRYSIFDVLRLDELEVILKAVLENSTAKDRPAVEKARTLYRSCMNQSV  
IEKRGSQPLLDILEVVGWPVAMDRWNETVGLEWELERQLALMNSQFNRRVLIDLFIWND  
DQNSSRHIIYIDQPTLGMPSPREYFFNGGSSNRKVVREAYLQFMVSVATLLREDANLPRDSC  
VQEDMMQVLELETQLAKATVPQEERHDVIALYHRMGLEELQSQFGLKGFNWTLFIQTVLS  
SVKIKLLPDEEVVVGIPYQLQNLNIIDTYSARTIQNYLVWRLVLDRIGSLSQRFKDTRV  
NYRKALFGTMVEEVRWRECVGYVNSNMENAVGSLYVREAFPGDSKSMVRELIDKVRTVFF  
ETLDELGWMDEESKKKAQEKAMSTREQIGHPDYILEEMNRRLDEEYSNLFNSEDLYFENS  
LQNLKVGAAQORSLRKLREKVDPNLWIIGAAVVNAFYSPNRNQIVFPAGILQPPFFSKEQPQ  
ALNFGGIGMVIGHEITHGFDDNGRNFDKNGNMMDWWNSNFSTQHFREQSECMIYQYGNYSW  
DLADEQNVNGFNTLGENIADNGGVRQAYKAYLKWMAEGGKDQQLPGLDLTHEQLFFINYA  
QVWCGSYRPEFAIQSIKTDVHSPLKYRVLGSLQNLAAAFADTFHCARGTPMHPKERCVRW

5'-

ATGGGGGAAGTCCGAAGGCCAGTGGGGATGGTGGAGAGCGCCGGCCGTGCAGGGCAGAAG  
CGCCCCGGGGTTCCCTGGAGGGGGGGCTGCTGCTGCTGCTGCTGCTGGTGACCGCTGCCCTG  
GTGGCCTTGGGTGTCCTCTACGCCGACCGCAGAGGGATCCCAGAGGCCCAAGAGGTGAGC  
GAGGTCTGCACCACCCCTGGCTGCGTGATAGCAGCTGCCAGGATCCTCCAGAACATGGAC  
CCGACCACGGAACCGTGAGCACTTCTACCAGTTTGCAATGCGGAGGCTGGCTGCGGCGC  
CACGTGATCCCTGAGACCAACTCAAGATACAGCATCTTTGACGTCCTCCGCGACGAGCTG  
GAGGTCACTCCTCAAAGCGGTGCTGGAGAATTGCACTGCCAAGGACCGGCCCGCTGTGGAG

AAGGCCAGGACGCTGTACCGCTCCTGCATGAACCAGAGTGTGATAGAGAAGCGAGGCTCT  
CAGCCCCCTGCTGGACATCTTGGAGGTGGTGGGAGGCTGGCCGGTGGCGATGGACAGGTGG  
AACGAGACCGTAGGACTCGAGTGGGAGCTGGAGCGGCAGCTGGCGCTGATGAACTCACAG  
TTCAACAGGCGCGCTCCTCATCGACCTCTTCATCTGGAACGACGACCAGAACTCCAGCCGG  
CACATCATCTACATAGACCAGCCCACCTTGGGCATGCCCTCCCGAGAGTACTACTTCAAC  
GGCGGCAGCAACCGGAAGGTGCGGGAAGCCTACCTGCAGTTCATGGTGTGAGTGGCCACG  
TTGCTGCGGGAGGATGCAAACCTGCCCAGGGACAGCTGCCCTGGTGCAGGAGGACATGATG  
CAGGTGCTGGAGCTGGAGACACAGCTGGCCAAGGCCACGGTACCCCAGGAGGAGAGACAC  
GACGTCATCGCCTTGTACCACCGGATGGGACTGGAGGAGCTGCAAAGCCAGTTTGGCCTG  
AAGGGATTTAACTGGACTCTGTTTCATACAAACTGTGCTATCCTCTGTCAAATCAAGCTG  
CTGCCAGATGAGGAAGTGGTGGTCTATGGCATCCCCCTACCTGCAGAACCTTGAAAACATC  
ATCGACACCTACTCAGCCAGGACCATAACAGAACTACCTGGTCTGGCGCCTGGTGGTGGAC  
CGCATTGGTAGCCTAAGCCAGAGATTCAAGGACACACGAGTGAACCTACCGCAAGGCGCTG  
TTTGGCACAATGGTGGAGGAGGTGCGCTGGCGTGAATGTGTGGGCTACGTCAACAGCAAC  
ATGGAGAACGCCGTGGGCTCCCTCTACGTCAGGGAGGCGTTCCCTGGAGACAGCAAGAGC  
ATGGTCAGAGAACTCATTGACAAGGTGCGGACAGTGTTTGTGGAGACGCTGGACGAGCTG  
GGCTGGATGGACGAGGAGTCCAAGAAGAAGGCGCAGGAGAAGGCCATGAGCATCCGGGAG  
CAGATCGGGCACCCTGACTACATCCTGGAGGAGATGAACAGGCGCCTGGACGAGGAGTAC  
TCCAATCTGAACTTCTCAGAGGACCTGTACTTTGAGAACAGTCTGCAGAACCTCAAGGTG  
GGCGCCCAGCGGAGCCTCAGGAAGCTTCGGGAAAAGGTGGACCCAAATCTCTGGATCATC  
GGGGCGGCGGTGGTCAATGCGTTCTACTCCCCAAACCGAAACCAGATTGTATTCCCTGCC  
GGGATCCTCCAGCCCCCTTCTTCAGCAAGGAGCAGCCACAGGCCTTGAACCTTTGGAGGC  
ATTGGGATGGTGATCGGGCACGAGATCACGCACGGCTTTGACGACAATGGCCGGAACCTC  
GACAAGAATGGCAACATGATGGATTGGTGGAGTAACCTTCTCCACCCAGCACTTCCGGGAG  
CAGTCAGAGTGCATGATCTACCAGTACGGCAACTACTCCTGGGACCTGGCAGACGAACAG  
AACGTGAACGGATTCAACACCCTTGGGGAAAACATTGCTGACAACGGAGGGGTGCGGCAA  
GCCTATAAGGCCTACCTCAAGTGGATGGCAGAGGGTGGCAAGGACCAGCAGCTGCCCCGC  
CTGGATCTCACCCATGAGCAGCTCTTCTTCATCAACTACGCCCAGGTGTGGTGCGGGTCC  
TACCGGCCCCGAGTTCGCCATCCAATCCATCAAGACAGACGTCCACAGTCCCCTGAAGTAC  
AGGGTACTGGGGTCGCTGCAGAACCTGGCCGCCTTCGCAGACACGTTCCACTGTGCCCGG  
GGCACCCCCATGCACCCCAAGGAGCGATGCCGCGTGTGGTAG - 3'

**Table 6: IGS5-protein-2 ("IGS5PROT2") of SEQ ID NO:6**

MGKSEGPVGMVESAGRAGQKRPGFLEGGLLLLLLLLVTAALVALGVLYADRRGIPEAQEVS  
EVCTTPGCVIAAARILQNMDPTEPCDDFYQFACGGWLRHVIPETNSRYSIFDVLDEL  
EVILKAVLENSTAKDRPAVEKARTLYRSCMNQSVIEKRGSQLLDILEVGGWFPVAMDRW  
NETVGLEWELERQLALMNSQFNRRVLIDLFIWNDDQNSSRHIIYIDQPTLGMPsREYYFN  
GGSNRKVREAYLQFMVSVATLLREDANLPRDSCLVQEDMMQVLELETQLAKATVPQEEH  
DVIALYHRMGLEELQSQFGLKGFNWTLEFIQTVLSSVKIKLLPDEEVVVGIPYLQNLNI  
IDTYSARTIQNYLVWRLVLDRIQSLSQRFKDTRVNYRKALFGTMVEEVRWRECVGYVNSN  
MENAVGSLYVREAFPGDSKSMVRELIDKVRTVFVETLDELGWMDEESKKKAQEKAMSIRE  
QIGHPDYILEEMNRRLDEEYSNLNFSEDLYFENSLQNLKVGAQRSRLKRLREKVDPNLWII  
GAAVVNAFYSPNRNQIVFPAGILOPPFFSKEQPQALNFGGIGMVGHEITHGFDDNGRNF  
DKNGNMDWWSNFSQHFREQSECMIYQYGNYSWDLADEQNVNGFNTLGENIADNGGVRQ  
AYKAYLKWMAEGGKDQQLPGLDLTHEQLFFINYAQVWCGSYRPEFAIQSIKTDVHSPKLY  
RVLGSLQNLAAAFADTFHCARGTPMHPKERCVRW

5 All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

10 The following examples are only intended to further illustrate the invention, in more detail, and therefore these examples are not deemed to restrict the scope of the invention in any way.

15 **EXAMPLE 1. THE CLONING OF cDNA ENCODING A NOVEL MEMBER OF THE NEPRILYSIN NEP/ECE METALLOPROTEASE FAMILY.**

**Example 1a. Outline of homology cloning of cDNA coding sequence of IGS5.**

20 Metalloproteases of the M13 subfamily are involved in the metabolism of various neuronal and hormonal peptides. To date this subfamily comprises neprilysin (NEP), endothelin-converting enzyme-1 (ECE-1), ECE-2, Kell, Pax and XCE.



Inhibitors of NEP and ECE are being developed for therapeutical use for example in cardiology and gastroenterology. Since additional members of this family may be interesting drug targets, homology cloning was used to identify novel genes in the human genome.

- 5 Homology cloning of IGS5 was performed according to the general description supra and according to the experimental details further illustrated by the experimental section of the international patent application PCT/EP 00/11532 which is incorporated by reference herein. The procedure may be outlined as follows:

- 10 In the databank of expressed sequence tags (ESTs), sequences were detected which contained a small open reading frame that showed similarity to the C-terminal part of NEP/ECE-like metalloproteases. Based on these EST sequences and conserved peptide motifs of the NEP/ECE-like metalloproteases we used degenerate PCR to clone the complete cDNA sequence from human lung, heart and testis cDNA.

- 15 The cDNA sequence encoded a glycosylated protein, named IGS5 or alternatively human soluble endopeptidase (hSEP), displaying the characteristics of M13 family members. IGS5 showed high amino acid sequence identity to mouse SEP (78%), mouse, rat and human NEP (54%) and to human ECE-1 (38%). In analogy with the mouse SEP and SEP<sup>A</sup> splice variants we also detected two IGS5 splice forms which differed in a 78bp alternative exon. These splice forms encoded proteins of 753 and 779 residues, respectively. The longer form contains a putative proteolytic cleavage site located adjacent to the transmembrane anchor. The two splice variants may  
20 therefore represent a membrane bound and a soluble form of the IGS5 protein.

- Expression analysis using multiple tissue dot blot analysis and quantitative PCR revealed expression in a variety of human tissues, with the strongest signal observed in testis. IGS5 mRNA expression was also observed e.g. in prostate, small intestine, stomach, colon, kidney and brain. The two splice variants showed a distinct expression pattern.

- 25 The functional characterization confirmed that this enzyme is a genuine member of the neprilysin family, and possesses ECE activity.

**Example 1b. Alignment of IGS5 with protein sequences of members of the NEP/ECE metalloprotease family.**

- 30 For the IGS5 Sequence originally cloned (see example 1), homology searches of up to date protein databanks and translated DNA databanks were executed using the BLAST algorithm (Altschul S.F. et al. [1997], Nucleic Acids Res. 25:3389-3402). These searches showed that the originally obtained IGS5 protein was most similar (54-55% identities over  $\pm$  700 aligned residues) to mouse, rat and human neutral endopeptidase (SW:NEP\_MOUSE, accession n° Q61391;  
35 SW:NEP\_RAT, accession n° P07861 and SW:NEP\_HUMAN accession n° P08473). Thus, this alignment of the almost complete IGS5 protein sequence with the other members of the NEP/ECE

family shows the relation of IGS5 to metalloproteases in general, and in particular to the NEP and/or ECE metalloprotease families. From this structural alignment it is concluded that the IGS5 has the functionality of metalloproteases, which in turn are of interest in the context of several dysfunctions, disorders or diseases in animals and humans.

5

**Example 1c. Cloning of cDNA fragments containing the full length coding sequence of IGS5.**

10 In order to obtain additional IGS5 cDNA sequence another round of RT-PCR reactions was carried out on human lung RNA under the conditions described above using the IGS5 specific reverse primer. The resulting contig contained an open reading frame which started at an "ATG" initiation codon and encoded a protein which showed high similarity with the N-terminal sequence of the mouse SEP protein.

15 Assembly of the DNA sequences of all isolated clones showed the presence of two types of cDNA sequences, that differed by the presence or absence of the 78 bp segment, initially identified within genomic clone IGS5/S1. These two sequences likely originate from alternatively spliced RNA molecules. The longest transcript contains an open reading frame of 2337 nucleotides (encoding a protein of 779 residues) whereas the shorter transcript contains an open reading

20 frame of 2259 nucleotides (encoding a protein of 753 residues). The coding sequence and the protein sequence of the long form is referred to as IGS5DNA1 (shown in SEQ ID NO:3, 2340 bp including the stop codon tag) and IGS5PROT1 (SEQ ID NO:4) respectively, whereas the coding sequence and the protein sequence of the shorter form are referred to as IGS5DNA2 (shown in SEQ ID NO:5, 2262 bp including the stop codon tag) and IGS5PROT2 (SEQ ID NO:6)

25 respectively. Downstream of the postulated methionine initiation codon within IGS5DNA1 and IGS5DNA2 an additional in-frame methionine codon is present at codon position 10. Although it is opted for the first methionine codon as being the initiation codon some (or even exclusive) initiation of translation at codon position 10 cannot be excluded, since both methionines appear to be within an equally favorable "Kozak" initiation of translation context (Kozak M., Gene [1999]:

30 234: 187-208). Hydropathy analysis (Kyte J. et al., J. Mol. Biol. [1982] 157: 105-132; Klein P. et al., Biochim. Biophys. Acta [1985] 815: 468-476) of the IGS5PROT1 and IGS5PROT2 sequences showed the presence of a single transmembrane domain between residues 22 to 50. This indicates that IGS5PROT1 and IGS5PROT2 are type II integral membrane proteins and thus have a membrane topology similar to other members of the NEP/ECE protein family.

35

**Example 1c. Alignment of IGS5 protein sequences of example 1c with protein sequences of members of the NEP/ECE metalloprotease family.**

For the IGS5 sequence cloned in example 1c, homology searches of up to date protein databanks and translated DNA databanks were executed using the BLAST algorithm (Altschul S.F. et al, Nucleic Acids Res. [1997] 25:3389-3402). These searches showed that IGS5PROT1 was most similar (76% identities over 778 aligned residues) to mouse SEP (GenBank accession n° AF157105) and also showed 54-55 % identities over 896 aligned residues to mouse, rat and human neutral endopeptidases (SW:NEP\_MOUSE, accession n° Q61391; SW:NEP\_RAT, accession n° P07861; SW:NEP\_HUMAN, accession n° P08473). Homology searches of IGS5PROT2 showed that this sequence was most similar (78% identities over 752 aligned residues) to mouse SEP<sup>d</sup> (GenBank accession no AF157106). In analogy with the mouse SEP and SEP<sup>d</sup> proteins it is to be expected that IGS5PROT1 and IGS5PROT2 represent the soluble and membrane-bound forms of the IGS5 protein, respectively. This is corroborated by the presence of dibasic residues (KRK) encoded at the 3' end of the the alternatively spliced 78bp exon.

Thus, this alignment of the complete IGS5 protein sequence with the other members of the NEP/ECE family shows the relation of IGS5 to NEP/ECE metalloproteases in general, and in particular to the SEP and NEP family members. From this structural alignment it is concluded that the IGS5 protein has the functionality of metalloproteases, which in turn are of interest in the context of several dysfunctions, disorders or diseases in animals and humans.

## **EXAMPLE 2. EXPRESSION AND PURIFICATION OF THE SOLUBLE HIS-TAGGED ECTODOMAIN OF HUMAN IGS5.**

The aim of the experiment was to produce soluble IGS5 protein using the baculoviral expression system. A recombinant baculovirus was constructed that expressed the His<sub>8</sub>-tagged IGS5 ectodomain upon infection of the Sf9 cell-line. Soluble IGS5 protein was then purified from the culture supernatant in a two step procedure involving lentil-lectin and Zn-IMAC chromatography, as was done in the state of the art for His<sub>8</sub>-ECE-1.

### **Example 2a. Experimental procedures.**

#### **Kinetic expression analysis.**

519 cells (IGCL 83.0), exponentially growing in suspension in Spinner flasks at 27°C in TC100 medium (JRH Biosciences cat n° 56941), supplemented with 10% inactivated Foetal Calf Serum (Gibco BRL cat n° 10 084 168), were collected by low speed centrifugation and seeded at  $5 \cdot 10^5$  cells/Fk (25 cm<sup>2</sup>) in serum-free TC100 medium. Candidate recombinant viral clones were added at a multiplicity of infection (MOI) of 3 pfu/cell and cell/virus cultures were subsequently incubated at 27°C. Cells and conditioned medium (CM) were harvested at 24, 48 and 72 h post infection by 2 consecutive low speed centrifugations. Samples were analysed by SDS PAGE gel electrophoresis

and Western blotting.

#### Deglycosylation study.

5 Samples were supplemented with SDS to a final concentration of 1% and incubated at 95°C for 5 min. After addition of 1 volume of the 2x Incubation buffer (250mM phosphate buffer, 50 mM EDTA, 5% N-octylglycoside, 1% 2-mercaptoethanol) and an additional 5 min incubation time at 95°C, the sample was cooled to 37°C. 1 U of N-glycosidase F (Boehringer Mannheim, cat n° 1 365 177) was added and after overnight incubation at 37°C, the sample was reduced with 100mM DTT (final concentration).

10

#### Preparative production.

519 cells (IGCL 83-2) exponentially growing in suspension in spinner flasks at 27°C in TC100 medium (JRH Biosciences, cat n° 56941) supplemented with 10% inactivated Foetal Calf Serum (Gibco BRL, cat n° 10 084 168) were collected by low speed centrifugation and resuspended at 15 density of  $2 \cdot 10^6$  cells/ml in TC100 medium, supplemented with 0.013 TIU aprotinin/ml (Pentex). Recombinant virus IGBV73 was added to the cells at a multiplicity of infection (MOI) of 2.25 pfu/cell (in stead of MOI 3 due to the low titre of the primary virus bank). The cell/virus suspension was subsequently incubated at 27°C in glass roller bottles (3 x 500 ml / 1260 cm<sup>2</sup>) for 72 h. The CM (1.5 l) was then cleared from cells and cell debris by two consecutive low speed 20 centrifugations. Aliquots were taken for quality control by Western blot analysis and for the determination of endotoxin levels.

#### Example 2b. Results.

25

#### Kinetics of expression.

The kinetics of expression of three candidate recombinant viral clones were studied via Western blot analysis. Western blot revealed a clear band at approximately 81 kDa in the CM of all candidate clones, corresponding to the theoretical Mr of the mature protein (81.2 kDa). For cell 30 lysates, the SDS gel was overloaded so no conclusions could be drawn. Expression levels of all 3 clones peaked at 48 to 72 h post-infection. Clone 2 was selected for further amplification and was deposited as IGBV73. Optimal harvest time was set at 72 h post infection.

#### Deglycosylation study.

35 The soluble IGS5 protein sequence contains 8 potential N-glycosylation sites. Since the purification protocol involves binding of the sugar residues on a lentil-lectin column samples of CM and cell lysates, harvested at 72h post infection were used for a deglycosylation study with N-glycosidase F, to check whether the recombinant soluble His<sub>6</sub>IGS5 protein is indeed expressed as

a glycosylated protein.

Western blot analysis of N-glycosidase F treated CM samples and non-treated controls show a shift in Mr when samples are deglycosylated, demonstrating that the soluble human His-tagged IGS5 is expressed as a glycosylated protein. In the non treated cell lysates, 3 protein bands of approximately 80 to 82 kDa can be observed. Upon N-glycosidase F treatment, 1 band of approximately 80 kDa remains visible, corresponding to the lowest MW band of the non treated samples.

#### 10 Preparative production.

1.5 liter of CM was harvested from IGBV73 infected Sf9 insect cells 72 h post infection.

Endotoxin content was determined to be 0.0847 EU/ml CM. Western blot analysis revealed a clear band at approximately 81 kDa in the CM, corresponding to the MW of the mature soluble His-tagged IGS5. When compared to cell lysate samples, which showed 3 protein bands, the CM protein band corresponds to the weaker middle Mr band, present in the cells.

### EXAMPLE 3. PURIFICATION OF IGS5.

#### 20 Example 3a. Experimental procedures.

##### Sample pretreatment.

1 tablet of EDTA free complete (EFC; Roche Biochemicals, cat n° 1873580) was added to 300 ml cleared Baculo CM. HEPEs, glycerol and Tween 20 were added to a final concentration of resp. 20 mM, 5% (v/v) and 0.005 % (w/v). The pH of the CM was adjusted to 7.4 and the sample was filtrated Durapore Membrane Filters 0.2 µ GV). All purification steps were performed at 4°C.

##### Lentil Lectin Chromatography.

The baculo sample was loaded overnight at 0.3 ml/min on a 5 ml Lentil Lectin Sepharose resin in a C10/10 column ( Pharmacia), which had been equilibrated in buffer A (20 mM Hapes, 150 mM NaCl, 5% glycerol, 0.005% Tween 20) supplemented with 1 tablet EFC / 500 ml. The column was washed with equilibration buffer until the absorbance at 280 nm reached baseline level and the bound proteins were eluted by applying buffer A containing 0.5 M alpha-methylpyrannoside. The column was regenerated by applying 100 mM acetate, 500 mM NaCl, pH 5.0. The elution and regeneration liquids were collected manually and the pools were analyzed by SDS-PAGE on 12.5 % Phast gels (Pharmacia) and silver staining. Prestained markers (Gibco) were included as relative molecular weight (Mr) standard.

**Immobilized metal affinity chromatography (IMAC) and dialysis.**

1 ml Chelating HiTrap (Pharmacia) was loaded with zinc ions as described by the manufacturer and equilibrated with buffer B (20 mM Hepes, 100 mM NaCl, 5% glycerol, 0.005% (w/v) Tween 20, pH 7.2). Lentil elution pools 1 and 2 were loaded separately at 0.5 ml/min on the HiTrap column (IMAC run A and IMAC run B). A blank run was included to compare the chromatographic absorbance profile. The column was washed with buffer B till baseline level and bound proteins were eluted by applying an imidazole step gradient (20, 50, 100 and 200 mM) in buffer B. Fractions were collected manually. The IMAC column was regenerated by applying 20 mM Hepes, 50 mM EDTA, 500 mM NaCl, pH 7.2. Elution and regeneration pools were analyzed by SDS-PAGE (12.5% Phast gels, Pharmacia) and silver staining. The 200 mM imidazole pool was transferred to a slide a-lyzer-cassette (MWCO 10.000, Pierce) and dialyzed overnight against buffer B (130 fold excess, no buffer refreshment).

**Protein quantification.**

The amount of soluble IGSS in the dialyzed pool was determined with the micro-BCA method (Pierce). BSA was included as reference.

**Protein Characterization.**

The dialyzed baculo IGSS was biochemically characterized by (1) SDS-PAGE under reducing and non reducing conditions and (2) Western blot with an anti His-tag mAb (21E1B4, IG) followed by incubation with alkaline phosphatase labeled rabbit anti-mouse Ig (Dako) and detection with NBT/BCIP staining. The glycosilation status of the soluble IGSS was verified by PGNase F treatment (Biorad).

**Example 3b. Results.**

The Lentil elution profile with 0.5 M alfa-methylpyrannoside resulted in a tailing peak. The flow through, washes and the elution pools were analyzed by SDS-PAGE and silver staining. The major amount of proteins were retrieved in the flow through and an IGSS-candidate band with a Mr of about 85.000 was observed in elution pools 1 to 3. Western blot analysis of the lentil chromatography with the anti-His tag mAb showed that the soluble hIGS5 protein (Mr ~ 85.000) is quantitatively bound to the Lentil Lectin resin and that the His-tagged protein is recovered over the whole elution peak, but mainly in pools 1 and 2. The Lentil Lectin elution pools 1 and 2 were further processed on the zinc-IMAC column (runs A and B). The bound proteins were eluted by an imidazole step gradient. SDS-PAGE analysis and silver staining showed that the bulk of contaminating proteins were eluted by applying the 20 mM and 50 mM imidazole step. The hIGS5 protein was retrieved in the 100 mM and 200 mM imidazole elution steps. The 100 mM imidazole

elution, which contains maximum 10% of the hIGS5 in the eluate, is still contaminated with a protein with a Mr of > 115.000. The IGS5 band in the 100 mM is also a doublet band. It remains to be verified whether the faint upper band, which represents less than 10% of the doublet, is a residual baculo contaminant or an IGS5 isoform or whether the lower (intense) band is a carboxy-terminal degradation product. The 85 kDa band in the 200 mM imidazole pool is a single band on the SDS-PAGE, which reacts with the anti his-tag mAb.

Silver staining did not reveal any difference in purity between the hIGS5 material obtained from IMAC run A and run B. This indicates that pool 1 and pool 2 of the Lectin eluate can be pooled in the future and be processed simultaneously in a single run on the zinc-IMAC column.

An identical band pattern was observed on Coomassie stained SDS-PAGE gels run under reducing and non reducing conditions, indicating that the purified hIGS5 does not contain disulfide based oligomers. Treatment with PGNase F reduced the Mr on SDS-PAGE with about 5 kDa. This minimal shift of the baculo expressed hIGS5 after treatment with endoglycosidase differs strongly from the migration shift which was observed for CHO expressed mouse SEP (Ikeda et al., JBC, 274, 32469, 1999).

Starting from 300 ml of baculo CM, 340 µg of over 95% pure His-tagged hIGS5 ectodomain was obtained by the 2 step purification procedure (i.e. a yield of about 1 mg/l). The purified product was then used in the enzyme inhibition assays as indicated in the following examples.

#### EXAMPLE 4. IGS5 ENZYME INHIBITION ASSAY.

The enzymatic activity of IGS5 polypeptides of the invention was tested with regard to the metabolism of biologically active peptides. In particular it was tested whether these IGS5 polypeptides may act on a variety of vasoactive peptides known in the state of the art e.g. such like atrial natriuretic peptide (ANP), bradykinin, big-endothelin (big-ET-1), endothelin (ET-1), substance P and angiotensin-1. In the context of the present invention in particular it was tested whether the IGS5 ectodomain, which is a novel human metalloprotease, hydrolyzes said vasoactive peptides. For comparison the assay was also performed for a known member of the metalloprotease family which was described earlier as soluble secreted endopeptidase (SEP) by Emoto et al. (J. Biol. Chem., Vol. 274 (1999); pp. 32469-32477). Furthermore, it was tested whether the activity of IGS5 to convert a big-ET-1 analogue (the so-called 17 aa big-ET-1) may be inhibited by reference compounds that are used to determine the inhibition properties with regard to enzymes having ECE and/or NEP-characteristics. Compounds used to test the inhibition of IGS5-activity on the big-ET-1 analogue were the compound phosphoramidon which inhibits

endopeptidases with NEP-characteristics, the compound thiorphan which selectively inhibits NEP, and the compound CGS-35066 which is a dualNEP/ECE inhibitor.

#### Example 4a. Materials.

- 5 Enzyme: IGS-5 (sol hu)(his)8; or: His6-tagged IGS5 ectodomain;  
stock solution: 53 µg/ml in 20 mM HEPES pH 7.2, 5% glycerol, 0.005% Tween20,  
100 mM NaCl, purity >99%; storage at 4 °C.  
working solution: stock solution diluted with assay buffer to 10 µg/ml.
- Substrate: Mca-Asp-Ile-Ala-Trp-Phe-Dpa-Thr-Pro-Glu-His-Val-Val-Pro-  
10 Tyr-Gly-Leu-Gly-COOH;  
Fluorescence-quenched big-ET-1 analogue;  
Mca = 7-Methoxycoumarin-4-yl;  
Dpa = 3-[2,4-Dinitrophenyl]-L-2,3-diaminopropionyl;  
stock solution: 100 µM in assay buffer; storage at -20 °C.
- 15 (commercially available from supplier: Polypeptide Laboratories,  
Wolfenbützel, Germany)
- Assay buffer: 100 mM Tris pH 7.0, 250 mM NaCl.

All test compounds were dissolved in DMSO at 10 mM and were further diluted with assay buffer.

20

#### Example 4b. Assay Procedure.

A quantity of 70 µl of the assay buffer, of 10 µl enzyme working solution and of 10 µl test compound solution were mixed in an Eppendorf vial and preincubated at 37 °C for 15 minutes. Then, 10 µl substrate stock solution was added and the reaction mixture was incubated at 37 °C  
25 for 60 minutes to allow for enzymatic hydrolysis. Subsequently the enzymatic reaction was terminated by heating at 95 °C for 5 minutes. After centrifugation (Heraeus Biofuge B, 3 min) the supernatant was subjected to HPLC analysis.

#### Example 4c. HPLC Procedure.

- 30 In order to separate the remaining substrate from the cleavage products reversed phase HPLC technique was used with a CC 125/4 Nucleosil 300/5 C<sub>18</sub> RP column and a CC 8/4 Nucleosil 100/5 C<sub>18</sub> precolumn (commercially available from Macherey-Nagel, Düren, Germany). Thus, 60 µl of the reaction samples obtained in Example 7b were injected into the HPLC, and the column was eluted at a flow rate of 1 ml/min by applying the following gradient and solutions:

- 35 Solution A: 100% H<sub>2</sub>O + 0.5 M H<sub>3</sub>PO<sub>4</sub>, pH = 2.0  
Solution B: 100% acetonitrile + 0.5M H<sub>3</sub>PO<sub>4</sub>



	0-2 min	20%	B
	2-6 min	20-60%	B
	6-8 min	60%	B
	8-10 min	60-90%	B
5	10-13 min	90%	B
	13-15 min	90-100%	B

Peptides were detected by absorbance at 214 nm and by fluorescence with an excitation wavelength of 328 nm and an emission wavelength of 393 nm.

10

**Example 4d. Calculations.**

The increasing fluorescence signal of the HPLC-peak of the peptide with the unquenched Mca-fluorophor after hydrolysis was taken as the basis for any calculation.

15

This signal was compared for the samples with and without inhibitor and % inhibition was calculated on basis of the respective peak areas.

$$\% \text{ Inhib} = 100 \cdot (1 - A_{\text{inhib}} / A_{\text{control}})$$

All samples were run in duplicate and mean values were used.

20

A standard inhibitor (10 nM and 100 nM Phosphoramidon) and a solvent control (0.1%) was added to each assay run.

**Example 4e. Results.**

25

With regard to the IGS5 polypeptides of the present invention the results of Example 4 show that these IGS5 metalloprotease polypeptides hydrolyze in vitro a variety of vasoactive peptides known in the state of the art. The results of the hydrolysis assay in comparison to the activity of SEP are shown in Table 7. From these results it is concluded that IGS5 may be particularly involved in the metabolism of said biologically vasoactive peptides.

30

**Table 7: Hydrolysis of vasoactive peptides by IGS5 polypeptides in comparison to SEP (soluble secreted end peptidase).**

Vasoactive Peptide	% Hydrolysis by IGS5 Polypeptide	% Hydrolysis by SEP (Emoto et al.)
	Conditions: 100 µg IGS5 polypeptide; 0.5 µM substrate; 2 h, 37 °C.	Conditions: 10 µg SEP; 0.5 µM substrate; 12 h, 37 °C.
ANP	5 (80*)	> 95
Bradykinin	100 (62**)	> 95
ET-1	< 30	92
Substance P	100	> 95
Angiotensin 1	15***	> 95
17 aa big-ET-1****	41	n.d.

\* 500 µg IGS5 polypeptide

\*\* 10 µg IGS5 polypeptide

5 \*\*\* degradation of angiotensin I does not result in formation of angiotensin II

\*\*\*\* 17 aa big-ET-1 was used as an analog of the natural big-ET-1; hydrolyzing activity of IGS5 was also detected using the natural big-ET-1, but could not be quantified due to difficulties with the HPLC-detection

#### EXAMPLE 5. NEP ENZYME INHIBITION ASSAY.

Neutral endopeptidase (E.C. 3.4.24.11) was prepared from pig kidney cortex according to the method of Gee et al. (Biochem J 1985 May 15;228(1):119-26) and purified as reported by Relton et al. (Biochem J 1983 Dec 1;215(3):519-23). For the enzyme inhibition assay 10 ng of the purified enzyme, 20 µM substrate (methionin-enkephalin) and various inhibitor concentrations were used. The assay buffer was 50 mM Tris-(hydroxymethyl)-aminomethan/HCl pH 7.4, the total assay volume was 100 µl. After a preincubation period of 5 min of enzyme and inhibitor, the substrate was added and subsequently a second incubation phase for enzyme induced substrate hydrolysis at 37°C for 30 min was initiated. The enzymatic reaction was stopped by heating at 95°C for 5 min. After centrifugation the supernatant was subjected to HPLC. The products of enzymatic substrate hydrolysis were separated from the native substrate by HPLC technique and the inhibitor potency calculated by comparing the peak areas of the products with the peak area of native substrate for both, the samples with and without inhibitor (control). Blanks without enzyme, controls without

inhibitor, samples with inhibitor solvent instead of the inhibitor and samples with a standard inhibitor were added to each assay run.

**Supplier of Materials:**

- 5 NEP: Dr. Philippe Crine, Univ. of Montreal, Canada  
Methionin-enkephalin: Sigma, Delsenhofen, Germany

**EXAMPLE 6. ECE ENZYME INHIBITION ASSAY.**

10

Recombinant human COOH-terminal His6-tagged endothelin converting enzyme-1 was expressed in Sf9-cells. Purification was performed by affinity chromatography. The enzyme inhibition assay comprised enzyme ( 2.8 µg), 5 µg substrate (moderately modified 17 amino acid truncated big endothelin-1), inhibitor at various final concentrations and 100 mM Tris-buffer (Tris-hydroxymethyl-aminomethan/HCl, pH 7.0 + 150 mM NaCl) in a final volume of 100 µl. Pre-incubation of enzyme with inhibitor for 15 min at 37°C was performed before substrate addition and incubation (60 min at 37°C) for enzymatic hydrolysis. The enzymatic reaction was stopped by heating at 95°C for 5 min. After centrifugation the supernatant was subjected to HPLC for separation of enzymatic hydrolysis products from undegraded substrate. % inhibition was calculated on the basis of peak areas for products and uncleaved substrate for the inhibited reaction in comparison to the control (without inhibitor). Blanks without enzyme, controls without inhibitor, samples with inhibitor solvent instead of the inhibitor and samples with a standard inhibitor were added to each assay run.

15

20

**Supplier of Materials:**

- 25 ECE: Innogenetics, Ghent, Belgium  
ECE substrate: Polypeptide, Wolfenbüttel, Germany

30

**EXAMPLE 7. BIOCHEMICAL PROFILE OF NEP/IGS5-INHIBITORY COMPOUNDS IN COMPARISON TO REFERENCE COMPOUNDS.**

35

In order to characterize and evaluate the pharmacological enzymatic properties of IGS5 for the purpose of the present invention a human IGS5 protein was generated by using an insect cell line as the expression system as described in the examples supra, and a variety of potential substrates of the IGS5 protein were tested. According to the results of example 4 IGS5 was found to efficiently cleave big-ET-1, ANP, and bradykinin, thus confirming that this novel protein is a genuine metalloprotease with a broad substrate specificity, which is a common feature of metalloproteases and which feature has been reported for NEP, ECE-1 and also ACE. It should

also be noted that according to the findings of the present invention the proteolysis of big-ET-1 by IGS5 surprisingly results in the correct formation of ET-1, e.g. big-ET-1 is correctly cleaved between amino acids Trp21 and Val22.

5 Furthermore, the potency of metalloprotease inhibitor compounds and reference compounds to suppress the conversion of big-ET to ET-1 was examined as described in this example 7, using a labeled fluorescent big-ET-1 analogue. The procedures applied in the enzyme assays are described above in the examples 4 with regard to IGS5, in example 5 with regard to NEP, and in example 6 with regard to ECE.

10

The results of the enzyme inhibition tests are summarized in Table 8. It is of interest that phosphoramidon that is known to inhibit the conversion of big-ET to ET-1 in vivo, also inhibits IGS5 with high potency in the biochemical assay used in the present invention, and surprisingly that the inhibition of IGS5 by phosphoramidon is actually considerably higher than ECE-1. In contrast, the selective NEP inhibitor thiorphan as well as the selective ECE-1 inhibitor SM-19712 (4-chloro-N-[[[(4-cyano-3-methyl-1-phenyl-1H-pyrazol-5-yl)amino]carbonyl] benzenesulfonamide, monosodium salt; Umekawa K, Hasegawa H, Tsutsumi Y, Sato K, Matsumura Y, Ohashi N., J Pharmacol 2000 Sep;84(1):7-15; Discovery Research Laboratories I, Research Center, Sumitomo Pharmaceuticals Co, Ltd, Osaka, Japan) do not affect the activity of IGS5 (Table 8).

20

**Table 8: Biochemical profile of NEP/IGS5-inhibitory compounds and reference compounds**

Compound	IGS5 IC <sub>50</sub> (nM)	NEP IC <sub>50</sub> (nM)	ECE IC <sub>50</sub> (nM)
Phosphoramidon	18	2.0	300
CGS-35066	1300	42	5
Thiorphan	> 1000	2.4	> 10000
SM-19712	> 10000	> 1000	11.7
compound Ia-2 active drug	1.2	2.9	1000
compound Ia-2 active drug	2.9	1.7	3279
compound Ia-2 active drug	2.8	4	374

The results of the present invention are very surprising, in particular in view of the following facts. Since endothelin converting enzyme-1 (ECE-1) was cloned in 1994, this enzyme has become generally accepted as the endopeptidase responsible for the physiological conversion of big-ET-1 to ET-1. However, despite the fact that ECE-1 has the ability to cleave big-ET-1, more recent reports raise doubts as to whether ECE-1 is the only physiologically relevant endothelin converting enzyme, or at least argue that additional enzymes must be involved in the production of ET-1. The IGS5 protein was discovered to be a metalloprotease with high similarity to NEP (54% identity) and a somewhat lower similarity to ECE-1 (39% identity). To characterize the enzymatic properties of this novel metalloprotease by the experiments under the example 4 it was looked for potential substrates and found that substance P, bradykinin, big-ET-1 and less efficiently ANP, angiotensin I and ET-1 are cleaved by IGS5. The enzymatic activity of IGS5 has an optimum at neutral pH (7.0-7.5). It should be noted that the proteolysis of big-ET-1 by IGS5 results in correct formation of ET-1.

By the experiments under the present example 7 the potency of metalloprotease inhibitors to suppress the conversion of big-ET to ET-1 by IGS5 was examined, using a labelled fluorescent big-ET-1 analogue as a substrate. It is of interest that phosphoramidon that is known to inhibit the conversion of big-ET to ET-1 in vivo, also inhibits IGS5 with high potency ( $IC_{50} = 18 \text{ nM}$ ) in our biochemical assay (considerably higher than ECE-1). In contrast, the selective NEP inhibitor thiorphan as well as a selective ECE-1 inhibitor SM-19712 from Sumitomo do not affect the activity of IGS5.

Thus compounds with formula I, in particular with formula Ia or Ib, respectively, surprisingly showed combined or concurrent NEP/IGS5-inhibitory activity. As representative examples of compounds of the formula Ia compound Ia-2 and of the formula Ib compound Ib-8 were investigated in the experiments of the present example 7. Both compounds revealed to be very potent inhibitors ( $IC_{50} = 1.2$  and  $2.9 \text{ nM}$ ) of the newly identified IGS5 metalloprotease.

54

## SEQUENCE LISTING

5 <110> SOLVAY PHARMACEUTICALS GmbH

<120> Use of Compounds with Combined NEP/MP-Inhibitory Activity in the Preparation of Medicaments

10 <130> HA 00.17-EP

<140>

<141>

15 <160> 6

<170> PatentIn Ver. 2.1

<210> 1

20 <211> 2076

<212> DNA

<213> Homo sapiens

<220>

25 <221> CDS

<222> (1)..(2073)

<400> 1

30 tgc acc acc cct ggc tgc gtg ata gca gct gcc agg atc ctc cag aac 48  
Cys Thr Thr Pro Gly Cys Val Ile Ala Ala Ala Arg Ile Leu Gln Asn  
1 5 10 15

atg gac ccg acc acg gaa cgg tgt gac gac ttc tac cag ttt gca tgc 96  
Met Asp Pro Thr Thr Glu Pro Cys Asp Asp Phe Tyr Gln Phe Ala Cys  
20 25 30

35 gga ggc tgg ctg cgg cgc cac gtg atc cct gag acc aac tca aga tac 144  
Gly Gly Trp Leu Arg Arg His Val Ile Pro Glu Thr Asn Ser Arg Tyr  
35 40 45

40 agc atc ttt gac gtc ctc cgc gac gag ctg gag gtc atc ctc aaa gcg 192  
Ser Ile Phe Asp Val Leu Arg Asp Glu Leu Glu Val Ile Leu Lys Ala  
50 55 60

45 gtg ctg gag aat tcg act gcc aag gac cgg ccg gct gtg gag aag gcc 240  
Val Leu Glu Asn Ser Thr Ala Lys Asp Arg Pro Ala Val Glu Lys Ala  
65 70 75 80

50 agg acg ctg tac cgc tcc tgc atg aac cag agt gtg ata gag aag cga 288  
Arg Thr Leu Tyr Arg Ser Cys Met Asn Gln Ser Val Ile Glu Lys Arg  
85 90 95

55 ggc tct cag ccc ctg ctg gac atc ttg gag gtg gtg gga ggc tgg ccg 336  
Gly Ser Gln Pro Leu Leu Asp Ile Leu Glu Val Val Gly Gly Trp Pro  
100 105 110

gtg gcg atg gac agg tgg aac gag acc gta gga ctc gag tgg gag ctg 384  
Val Ala Met Asp Arg Trp Asn Glu Thr Val Gly Leu Glu Trp Glu Leu  
115 120 125

60 gag cgg cag ctg gcg ctg atg aac tca cag ttc aac agg cgc gtc ctc 432  
Glu Arg Gln Leu Ala Leu Met Asn Ser Gln Phe Asn Arg Arg Val Leu  
130 135 140

## 55

	atc gac ctc ttc atc tgg aac gac gac cag aac tcc agc cgg cac atc	480
	Ile Asp Leu Phe Ile Trp Asn Asp Asp Gln Asn Ser Ser Arg His Ile	
	145 150 155 160	
5	atc tac ata gac cag ccc acc ttg ggc atg ccc tcc cga gag tac tac	528
	Ile Tyr Ile Asp Gln Pro Thr Leu Gly Met Pro Ser Arg Glu Tyr Tyr	
	165 170 175	
10	ttc aac ggc ggc agc aac cgg aag gtg cgg gaa gcc tac ctg cag ttc	576
	Phe Asn Gly Gly Ser Asn Arg Lys Val Arg Glu Ala Tyr Leu Gln Phe	
	180 185 190	
15	atg gtg tca gtg gcc acg ttg ctg cgg gag gat gca aac ctg ccc agg	624
	Met Val Ser Val Ala Thr Leu Leu Arg Glu Asp Ala Asn Leu Pro Arg	
	195 200 205	
20	gac agc tgc ctg gtg cag gag gac atg atg cag gtg ctg gag ctg gag	672
	Asp Ser Cys Leu Val Gln Glu Asp Met Met Gln Val Leu Glu Leu Glu	
	210 215 220	
25	aca cag ctg gcc aag gcc acg gta ccc cag gag gag aga cac gac gtc	720
	Thr Gln Leu Ala Lys Ala Thr Val Pro Gln Glu Glu Arg His Asp Val	
	225 230 235 240	
30	atc gcc ttg tac cac cgg atg gga ctg gag gag ctg caa agc cag ttt	768
	Ile Ala Leu Tyr His Arg Met Gly Leu Glu Glu Leu Gln Ser Gln Phe	
	245 250 255	
35	ggc ctg aag gga ttt aac tgg act ctg ttc ata caa act gtg cta tcc	816
	Gly Leu Lys Gly Phe Asn Trp Thr Leu Phe Ile Gln Thr Val Leu Ser	
	260 265 270	
40	tct gtc aaa atc aag ctg ctg cca gat gag gaa gtg gtg gtc tat ggc	864
	Ser Val Lys Ile Lys Leu Leu Pro Asp Glu Glu Val Val Val Tyr Gly	
	275 280 285	
45	atc ccc tac ctg cag aac ctt gaa aac atc atc gac acc tac tca gcc	912
	Ile Pro Tyr Leu Gln Asn Leu Glu Asn Ile Ile Asp Thr Tyr Ser Ala	
	290 295 300	
50	agg acc ata cag aac tac ctg gtc tgg cgc ctg gtg ctg gac cgc att	960
	Arg Thr Ile Gln Asn Tyr Leu Val Trp Arg Leu Val Leu Asp Arg Ile	
	305 310 315 320	
55	ggt agc cta agc cag aga ttc aag gac aca cga gtg aac tac cgc aag	1008
	Gly Ser Leu Ser Gln Arg Phe Lys Asp Thr Arg Val Asn Tyr Arg Lys	
	325 330 335	
60	gcg ctg ttt ggc aca atg gtg gag gag gtg cgc tgg cgt gaa tgt gtg	1056
	Ala Leu Phe Gly Thr Met Val Glu Glu Val Arg Trp Arg Glu Cys Val	
	340 345 350	
65	ggc tac gtc aac agc aac atg gag aac gcc gtg ggc tcc cto tac gtc	1104
	Gly Tyr Val Asn Ser Asn Met Glu Asn Ala Val Gly Ser Leu Tyr Val	
	355 360 365	
70	agg gag gcg ttc cct gga gac agc aag agc atg gtc aga gaa ctc att	1152
	Arg Glu Ala Phe Pro Gly Asp Ser Lys Ser Met Val Arg Glu Leu Ile	
	370 375 380	
75	gac aag gtg cgg aca gtg ttt gtg gag acg ctg gac gag ctg ggc tgg	1200
	Asp Lys Val Arg Thr Val Phe Val Glu Thr Leu Asp Glu Leu Gly Trp	
	385 390 395 400	

56

	atg gac gag gag tcc aag aag aag gcg cag gag aag gcc atg agc atc	1248
	Met Asp Glu Glu Ser Lys Lys Lys Ala Gln Glu Lys Ala Met Ser Ile	
	405 410 415	
5	cgg gag cag atc ggg cac cct gac tac atc ctg gag gag atg aac agg	1296
	Arg Glu Gln Ile Gly His Pro Asp Tyr Ile Leu Glu Glu Met Asn Arg	
	420 425 430	
10	cgc ctg gac gag gag tac tcc aat ctg aac ttc tca gag gac ctg tac	1344
	Arg Leu Asp Glu Glu Tyr Ser Asn Leu Asn Phe Ser Glu Asp Leu Tyr	
	435 440 445	
15	ttt gag aac agt ctg cag aac ctc aag gtg ggc gcc cag cgg agc ctc	1392
	Phe Glu Asn Ser Leu Gln Asn Leu Lys Val Gly Ala Gln Arg Ser Leu	
	450 455 460	
20	agg aag ctt cgg gaa aag gtg gac cca aat ctc tgg atc atc ggg gcg	1440
	Arg Lys Leu Arg Glu Lys Val Asp Pro Asn Leu Trp Ile Ile Gly Ala	
	465 470 475 480	
25	gcg gtg gtc aat gcg ttc tac tcc cca aac cga aac cag att gta ttc	1488
	Ala Val Val Asn Ala Phe Tyr Ser Pro Asn Arg Asn Gln Ile Val Phe	
	485 490 495	
30	cct gcc ggg atc ctc cag ccc ccc ttc ttc agc aag gag cag cca cag	1536
	Pro Ala Gly Ile Leu Gln Pro Pro Phe Phe Ser Lys Glu Gln Pro Gln	
	500 505 510	
35	gcc ttg aac ttt gga ggc att ggg atg gtg atc ggg cac gag atc acg	1584
	Ala Leu Asn Phe Gly Gly Ile Gly Met Val Ile Gly His Glu Ile Thr	
	515 520 525	
40	cac ggc ttt gac gac aat ggc cgg aac ttc gac aag aat ggc aac atg	1632
	His Gly Phe Asp Asp Asn Gly Arg Asn Phe Asp Lys Asn Gly Asn Met	
	530 535 540	
45	atg gat tgg tgg agt aac ttc tcc acc cag cac ttc cgg gag cag tca	1680
	Met Asp Trp Trp Ser Asn Phe Ser Thr Gln His Phe Arg Glu Gln Ser	
	545 550 555 560	
50	gag tgc atg atc tac cag tac ggc aac tac tcc tgg gac ctg gca gac	1728
	Glu Cys Met Ile Tyr Gln Tyr Gly Asn Tyr Ser Trp Asp Leu Ala Asp	
	565 570 575	
55	gaa cag aac gtg aac gga ttc aac acc ctt ggg gaa aac att gct gac	1776
	Glu Gln Asn Val Asn Gly Phe Asn Thr Leu Gly Glu Asn Ile Ala Asp	
	580 585 590	
60	aac gga ggg gtg cgg caa gcc tat aag gcc tac ctc aag tgg atg gca	1824
	Asn Gly Gly Val Arg Gln Ala Tyr Lys Ala Tyr Leu Lys Trp Met Ala	
	595 600 605	
65	gag ggt ggc aag gac cag cag ctg ccc ggc ctg gat ctc acc cat gag	1872
	Glu Gly Gly Lys Asp Gln Gln Leu Pro Gly Leu Asp Leu Thr His Glu	
	610 615 620	
70	cag ctc ttc ttc atc aac tac gcc cag gtg tgg tgc ggg tcc tac cgg	1920
	Gln Leu Phe Phe Ile Asn Tyr Ala Gln Val Trp Cys Gly Ser Tyr Arg	
	625 630 635 640	



## 57

ccc gag ttc gcc atc caa tcc atc aag aca gac gtc cac agt. ccc ctg 1968  
 Pro Glu Phe Ala Ile Gln Ser Ile Lys Thr Asp Val His Ser Pro Leu 655  
 645 650

5 aag tac agg gta ctg ggg tcc ctg cag aac ctg gcc gcc ttc gca gac 2016  
 Lys Tyr Arg Val Leu Gly Ser Leu Gln Asn Leu Ala Ala Phe Ala Asp 670  
 660 665

10 acg ttc cac tgt gcc cgg ggc acc ccc atg cac ccc aag gag cga tgc 2064  
 Thr Phe His Cys Ala Arg Gly Thr Pro Met His Pro Lys Glu Arg Cys 685  
 675 680

15 cgc gtg tgg tag 2076  
 Arg Val Trp 690

20 <210> 2  
 <211> 691  
 <212> PRT  
 <213> Homo sapiens

25 <400> 2  
 Cys Thr Thr Pro Gly Cys Val Ile Ala Ala Ala Arg Ile Leu Gln Asn 15  
 1 5 10

Met Asp Pro Thr Thr Glu Pro Cys Asp Asp Phe Tyr Gln Phe Ala Cys 30  
 20 25

30 Gly Gly Trp Leu Arg Arg His Val Ile Pro Glu Thr Asn Ser Arg Tyr 45  
 35

Ser Ile Phe Asp Val Leu Arg Asp Glu Leu Glu Val Ile Leu Lys Ala 60  
 50 55

35 Val Leu Glu Asn Ser Thr Ala Lys Asp Arg Pro Ala Val Glu Lys Ala 80  
 65 70 75

40 Arg Thr Leu Tyr Arg Ser Cys Met Asn Gln Ser Val Ile Glu Lys Arg 95  
 85 90

Gly Ser Gln Pro Leu Leu Asp Ile Leu Glu Val Val Gly Gly Trp Pro 110  
 100 105 110

45 Val Ala Met Asp Arg Trp Asn Glu Thr Val Gly Leu Glu Trp Glu Leu 125  
 115 120

Glu Arg Gln Leu Ala Leu Met Asn Ser Gln Phe Asn Arg Arg Val Leu 140  
 130 135

50 Ile Asp Leu Phe Ile Trp Asn Asp Asp Gln Asn Ser Ser Arg His Ile 160  
 145 150 155

55 Ile Tyr Ile Asp Gln Pro Thr Leu Gly Met Pro Ser Arg Glu Tyr Tyr 175  
 165 170

Phe Asn Gly Gly Ser Asn Arg Lys Val Arg Glu Ala Tyr Leu Gln Phe 190  
 180 185

60 Met Val Ser Val Ala Thr Leu Leu Arg Glu Asp Ala Asn Leu Pro Arg 205  
 195 200

58

	Asp	Ser	Cys	Leu	Val	Gln	Glu	Asp	Met	Met	Gln	Val	Leu	Glu	Leu	Glu	
	210						215					220					
5	Thr	Gln	Leu	Ala	Lys	Ala	Thr	Val	Pro	Gln	Glu	Glu	Arg	His	Asp	Val	
	225				230						235				240		
	Ile	Ala	Leu	Tyr	His	Arg	Met	Gly	Leu	Glu	Glu	Leu	Gln	Ser	Gln	Phe	
					245					250					255		
10	Gly	Leu	Lys	Gly	Phe	Asn	Trp	Thr	Leu	Phe	Ile	Gln	Thr	Val	Leu	Ser	
				260					265					270			
	Ser	Val	Lys	Ile	Lys	Leu	Leu	Pro	Asp	Glu	Glu	Val	Val	Val	Tyr	Gly	
			275					280					285				
15	Ile	Pro	Tyr	Leu	Gln	Asn	Leu	Glu	Asn	Ile	Ile	Asp	Thr	Tyr	Ser	Ala	
		290					295					300					
	Arg	Thr	Ile	Gln	Asn	Tyr	Leu	Val	Trp	Arg	Leu	Val	Leu	Asp	Arg	Ile	
20		305				310					315					320	
	Gly	Ser	Leu	Ser	Gln	Arg	Phe	Lys	Asp	Thr	Arg	Val	Asn	Tyr	Arg	Lys	
					325					330					335		
25	Ala	Leu	Phe	Gly	Thr	Met	Val	Glu	Glu	Val	Arg	Trp	Arg	Glu	Cys	Val	
				340					345					350			
	Gly	Tyr	Val	Asn	Ser	Asn	Met	Glu	Asn	Ala	Val	Gly	Ser	Leu	Tyr	Val	
			355					360					365				
30	Arg	Glu	Ala	Phe	Pro	Gly	Asp	Ser	Lys	Ser	Met	Val	Arg	Glu	Leu	Ile	
		370					375					380					
	Asp	Lys	Val	Arg	Thr	Val	Phe	Val	Glu	Thr	Leu	Asp	Glu	Leu	Gly	Trp	
35		385				390					395				400		
	Met	Asp	Glu	Glu	Ser	Lys	Lys	Lys	Ala	Gln	Glu	Lys	Ala	Met	Ser	Ile	
					405					410					415		
40	Arg	Glu	Gln	Ile	Gly	His	Pro	Asp	Tyr	Ile	Leu	Glu	Glu	Met	Asn	Arg	
				420					425					430			
	Arg	Leu	Asp	Glu	Glu	Tyr	Ser	Asn	Leu	Asn	Phe	Ser	Glu	Asp	Leu	Tyr	
			435					440					445				
45	Phe	Glu	Asn	Ser	Leu	Gln	Asn	Leu	Lys	Val	Gly	Ala	Gln	Arg	Ser	Leu	
		450					455					460					
	Arg	Lys	Leu	Arg	Glu	Lys	Val	Asp	Pro	Asn	Leu	Trp	Ile	Ile	Gly	Ala	
50		465				470					475					480	
	Ala	Val	Val	Asn	Ala	Phe	Tyr	Ser	Pro	Asn	Arg	Asn	Gln	Ile	Val	Phe	
				485					490					495			
55	Pro	Ala	Gly	Ile	Leu	Gln	Pro	Pro	Phe	Phe	Ser	Lys	Glu	Gln	Pro	Gln	
				500					505					510			
	Ala	Leu	Asn	Phe	Gly	Gly	Ile	Gly	Met	Val	Ile	Gly	His	Glu	Ile	Thr	
			515					520					525				
60	His	Gly	Phe	Asp	Asp	Asn	Gly	Arg	Asn	Phe	Asp	Lys	Asn	Gly	Asn	Met	
		530					535					540					

59

	Met	Asp	Trp	Trp	Ser	Asn	Phe	Ser	Thr	Gln	His	Phe	Arg	Glu	Gln	Ser	
	545					550					555					560	
5	Glu	Cys	Met	Ile	Tyr	Gln	Tyr	Gly	Asn	Tyr	Ser	Trp	Asp	Leu	Ala	Asp	
					565					570					575		
	Glu	Gln	Asn	Val	Asn	Gly	Phe	Asn	Thr	Leu	Gly	Glu	Asn	Ile	Ala	Asp	
				580					585					590			
10	Asn	Gly	Gly	Val	Arg	Gln	Ala	Tyr	Lys	Ala	Tyr	Leu	Lys	Trp	Met	Ala	
			595					600					605				
	Glu	Gly	Gly	Lys	Asp	Gln	Gln	Leu	Pro	Gly	Leu	Asp	Leu	Thr	His	Glu	
15		610				615						620					
	Gln	Leu	Phe	Phe	Ile	Asn	Tyr	Ala	Gln	Val	Trp	Cys	Gly	Ser	Tyr	Arg	
	625					630					635					640	
20	Pro	Glu	Phe	Ala	Ile	Gln	Ser	Ile	Lys	Thr	Asp	Val	His	Ser	Pro	Leu	
					645					650					655		
	Lys	Tyr	Arg	Val	Leu	Gly	Ser	Leu	Gln	Asn	Leu	Ala	Ala	Phe	Ala	Asp	
				660					665					670			
25	Thr	Phe	His	Cys	Ala	Arg	Gly	Thr	Pro	Met	His	Pro	Lys	Glu	Arg	Cys	
			675					680					685				
	Arg	Val	Trp														
30				690													
	<210> 3																
	<211> 2340																
	<212> DNA																
35	<213> Homo Sapiens																
	<220>																
	<221> CDS																
	<222> (1) .. (2337)																
40	<400> 3																
	atg	ggg	aag	tcc	gaa	ggc	ccc	gtg	ggg	atg	gtg	gag	agc	gct	ggc	cgt	48
	Met	Gly	Lys	Ser	Glu	Gly	Pro	Val	Gly	Met	Val	Glu	Ser	Ala	Gly	Arg	
	1				5					10					15		
45	gca	ggg	cag	aag	cgc	ccg	ggg	ttc	ctg	gag	ggg	ggg	ctg	ctg	ctg	ctg	96
	Ala	Gly	Gln	Lys	Arg	Pro	Gly	Phe	Leu	Glu	Gly	Gly	Leu	Leu	Leu	Leu	
				20				25					30				
50	ctg	ctg	ctg	gtg	acc	gct	gcc	ctg	gtg	gcc	ttg	ggt	gtc	ctc	tac	gcc	144
	Leu	Leu	Leu	Val	Thr	Ala	Ala	Leu	Val	Ala	Leu	Gly	Val	Leu	Tyr	Ala	
				35				40					45				
55	gac	cgc	aga	ggg	aag	cag	ctg	cca	cgc	ctt	gct	agc	cgg	ctg	tgc	ttc	192
	Asp	Arg	Arg	Gly	Lys	Gln	Leu	Pro	Arg	Leu	Ala	Ser	Arg	Leu	Cys	Phe	
	50					55						60					
60	tta	cag	gag	gag	agg	acc	ttt	gta	aaa	cga	aaa	ccc	cga	ggg	atc	cca	240
	Leu	Gln	Glu	Glu	Arg	Thr	Phe	Val	Lys	Arg	Lys	Pro	Arg	Gly	Ile	Pro	
	65					70					75					80	

60

		gag gcc caa gag gtg agc gag gtc tgc acc acc cct ggc tgc gtg ata	288
		Glu Ala Gln Glu Val Ser Glu Val Cys Thr Thr Pro Gly Cys Val Ile	
		85 90 95	
5		gca gct gcc agg atc ctc cag aac atg gac ccg acc acg gaa ccg tgt	336
		Ala Ala Ala Arg Ile Leu Gln Asn Met Asp Pro Thr Thr Glu Pro Cys	
		100 105 110	
10		gac gac ttc tac cag ttt gca tgc gga ggc tgg ctg cgg cgc cac gtg	384
		Asp Asp Phe Tyr Gln Phe Ala Cys Gly Gly Trp Leu Arg Arg His Val	
		115 120 125	
15		atc cct gag acc aac tca aga tac agc atc ttt gac gtc ctc cgc gac	432
		Ile Pro Glu Thr Asn Ser Arg Tyr Ser Ile Phe Asp Val Leu Arg Asp	
		130 135 140	
20		gag ctg gag gtc atc ctc aaa gcg gtg ctg gag aat tcg act gcc aag	480
		Glu Leu Glu Val Ile Leu Lys Ala Val Leu Glu Asn Ser Thr Ala Lys	
		145 150 155 160	
25		gac cgg ccg gct gtg gag aag gcc agg acg ctg tac cgc tcc tgc atg	528
		Asp Arg Pro Ala Val Glu Lys Ala Arg Thr Leu Tyr Arg Ser Cys Met	
		165 170 175	
30		aac cag agt gtg ata gag aag cga ggc tct cag ccc ctg ctg gac atc	576
		Asn Gln Ser Val Ile Glu Lys Arg Gly Ser Gln Pro Leu Leu Asp Ile	
		180 185 190	
35		ttg gag gtg gtg gga ggc tgg ccg gtg gcg atg gac agg tgg aac gag	624
		Leu Glu Val Val Gly Gly Trp Pro Val Ala Met Asp Arg Trp Asn Glu	
		195 200 205	
40		acc gta gga ctc gag tgg gag ctg gag cgg cag ctg gcg ctg atg aac	672
		Thr Val Gly Leu Glu Trp Glu Leu Glu Arg Gln Leu Ala Leu Met Asn	
		210 215 220	
45		tca cag ttc aac agg cgc gtc ctc atc gac ctc ttc atc tgg aac gac	720
		Ser Gln Phe Asn Arg Arg Val Leu Ile Asp Leu Phe Ile Trp Asn Asp	
		225 230 235 240	
50		gac cag aac tcc agc cgg cac atc atc tac ata gac cag ccc acc ttg	768
		Asp Gln Asn Ser Ser Arg His Ile Ile Tyr Ile Asp Gln Pro Thr Leu	
		245 250 255	
55		ggc atg ccc tcc cga gag tac tac ttc aac ggc ggc agc aac cgg aag	816
		Gly Met Pro Ser Arg Glu Tyr Tyr Phe Asn Gly Gly Ser Asn Arg Lys	
		260 265 270	
60		gtg cgg gaa gcc tac ctg cag ttc atg gtg tca gtg gcc acg ttg ctg	864
		Val Arg Glu Ala Tyr Leu Gln Phe Met Val Ser Val Ala Thr Leu Leu	
		275 280 285	
65		cgg gag gat gca aac ctg ccc agg gac agc tgc ctg gtg cag gag gac	912
		Arg Glu Asp Ala Asn Leu Pro Arg Asp Ser Cys Leu Val Gln Glu Asp	
		290 295 300	
70		atg atg cag gtg ctg gag ctg gag aca cag ctg gcc aag gcc acg gta	960
		Met Met Gln Val Leu Glu Leu Glu Thr Gln Leu Ala Lys Ala Thr Val	
		305 310 315 320	
75		ccc cag gag gag aga cac gac gtc atc gcc ttg tac cac cgg atg gga	1008
		Pro Gln Glu Glu Arg His Asp Val Ile Ala Leu Tyr His Arg Met Gly	
		325 330 335	

61

ctg gag gag ctg caa agc cag ttt ggc ctg aag gga ttt aac tgg act 1056  
 Leu Glu Glu Leu Gln Ser Gln Phe Gly Leu Lys Gly Phe Asn Trp Thr  
 340 345 350

5

ctg ttc ata caa act gtg cta tcc tct gtc aaa atc aag ctg ctg cca 1104  
 Leu Phe Ile Gln Thr Val Leu Ser Ser Val Lys Ile Lys Leu Leu Pro  
 355 360 365

10

gat gag gaa gtg gtg gtc tat ggc atc ccc tac ctg cag aac ctt gaa 1152  
 Asp Glu Glu Val Val Val Tyr Gly Ile Pro Tyr Leu Gln Asn Leu Glu  
 370 375 380

15

aac atc atc gac acc tac tca gcc agg acc ata cag aac tac ctg gtc 1200  
 Asn Ile Ile Asp Thr Tyr Ser Ala Arg Thr Ile Gln Asn Tyr Leu Val  
 385 390 395 400

20

tgg cgc ctg gtg ctg gac cgc att ggt agc cta agc cag aga ttc aag 1248  
 Trp Arg Leu Val Leu Asp Arg Ile Gly Ser Leu Ser Gln Arg Phe Lys  
 405 410 415

25

gac aca cga gtg aac tac cgc aag gcg ctg ttt ggc aca atg gtg gag 1296  
 Asp Thr Arg Val Asn Tyr Arg Lys Ala Leu Phe Gly Thr Met Val Glu  
 420 425 430

gag gtg cgc tgg cgt gaa tgt gtg ggc tac gtc aac agc aac atg gag 1344  
 Glu Val Arg Trp Arg Glu Cys Val Gly Tyr Val Asn Ser Asn Met Glu  
 435 440 445

30

aac gcc gtg ggc tcc ctc tac gtc agg gag gcg ttc cct gga gac agc 1392  
 Asn Ala Val Gly Ser Leu Tyr Val Arg Glu Ala Phe Pro Gly Asp Ser  
 450 455 460

35

aag agc atg gtc aga gaa ctc att gac aag gtg cgg aca gtg ttt gtg 1440  
 Lys Ser Met Val Arg Glu Leu Ile Asp Lys Val Arg Thr Val Phe Val  
 465 470 475 480

40

gag acg ctg gac gag ctg ggc tgg atg gac gag gag tcc aag aag aag 1488  
 Glu Thr Leu Asp Glu Leu Gly Trp Met Asp Glu Glu Ser Lys Lys Lys  
 485 490 495

45

gcg cag gag aag gcc atg agc atc cgg gag cag atc ggg cac cct gac 1536  
 Ala Gln Glu Lys Ala Met Ser Ile Arg Glu Gln Ile Gly His Pro Asp  
 500 505 510

tac atc ctg gag gag atg aac agg cgc ctg gac gag gag tac tcc aat 1584  
 Tyr Ile Leu Glu Glu Met Asn Arg Arg Leu Asp Glu Glu Tyr Ser Asn  
 515 520 525

50

ctg aac ttc tca gag gac ctg tac ttt gag aac agt ctg cag aac ctc 1632  
 Leu Asn Phe Ser Glu Asp Leu Tyr Phe Glu Asn Ser Leu Gln Asn Leu  
 530 535 540

55

aag gtg ggc gcc cag cgg agc ctc agg aag ctt cgg gaa aag gtg gac 1680  
 Lys Val Gly Ala Gln Arg Ser Leu Arg Lys Leu Arg Glu Lys Val Asp  
 545 550 555 560

60

cca aat ctc tgg atc atc ggg gcg gcg gtg gtc aat gcg ttc tac tcc 1728  
 Pro Asn Leu Trp Ile Ile Gly Ala Ala Val Val Asn Ala Phe Tyr Ser  
 565 570 575

## 62

	cca aac cga aac cag att gta ttc cct gcc ggg atc etc cag ccc ccc	1776
	Pro Asn Arg Asn Gln Ile Val Phe Pro Ala Gly Ile Leu Gln Pro Pro	
	580 585 590	
5	ttc ttc agc aag gag cag cca cag gcc ttg aac ttt gga ggc att ggg	1824
	Phe Phe Ser Lys Glu Gln Pro Gln Ala Leu Asn Phe Gly Gly Ile Gly	
	595 600 605	
10	atg gtg atc ggg cac gag atc acg cac ggc ttt gac gac aat ggc cgg	1872
	Met Val Ile Gly His Glu Ile Thr His Gly Phe Asp Asp Asn Gly Arg	
	610 615 620	
15	aac ttc gac aag aat ggc aac atg atg gat tgg tgg agt aac ttc tcc	1920
	Asn Phe Asp Lys Asn Gly Asn Met Met Asp Trp Trp Ser Asn Phe Ser	
	625 630 635 640	
20	acc cag cac ttc egg gag cag tca gag tgc atg atc tac cag tac ggc	1968
	Thr Gln His Phe Arg Glu Gln Ser Glu Cys Met Ile Tyr Gln Tyr Gly	
	645 650 655	
25	aac tac tcc tgg gac ctg gca gac gaa cag aac gtg aac gga ttc aac	2016
	Asn Tyr Ser Trp Asp Leu Ala Asp Glu Gln Asn Val Asn Gly Phe Asn	
	660 665 670	
30	acc ctt ggg gaa aac att gct gac aac gga ggg gtg egg caa gcc tat	2064
	Thr Leu Gly Glu Asn Ile Ala Asp Asn Gly Gly Val Arg Gln Ala Tyr	
	675 680 685	
35	aag gcc tac etc aag tgg atg gca gag ggt ggc aag gac cag cag ctg	2112
	Lys Ala Tyr Leu Lys Trp Met Ala Glu Gly Gly Lys Asp Gln Gln Leu	
	690 695 700	
40	ccc ggc ctg gat etc acc cat gag cag etc ttc ttc atc aac tac gcc	2160
	Pro Gly Leu Asp Leu Thr His Glu Gln Leu Phe Phe Ile Asn Tyr Ala	
	705 710 715 720	
45	cag gtg tgg tgc ggg tcc tac cgg ccc gag ttc gcc atc caa tcc atc	2208
	Gln Val Trp Cys Gly Ser Tyr Arg Pro Glu Phe Ala Ile Gln Ser Ile	
	725 730 735	
50	aag aca gac gtc cac agt ccc ctg aag tac agg gta ctg ggg tcc ctg	2256
	Lys Thr Asp Val His Ser Pro Leu Lys Tyr Arg Val Leu Gly Ser Leu	
	740 745 750	
55	cag aac ctg gcc gcc ttc gca gac acg ttc cac tgt gcc cgg ggc acc	2304
	Gln Asn Leu Ala Ala Phe Ala Asp Thr Phe His Cys Ala Arg Gly Thr	
	755 760 765	
60	ccc atg cac ccc aag gag cga tgc cgc gtg tgg tag	2340
	Pro Met His Pro Lys Glu Arg Cys Arg Val Trp	
	770 775	
55	<210> 4	
	<211> 779	
	<212> PRT	
	<213> Homo Sapiens	
60	<400> 4	
	Met Gly Lys Ser Glu Gly Pro Val Gly Met Val Glu Ser Ala Gly Arg	
	1 5 10 15	

63

Ala Gly Gln Lys Arg Pro Gly Phe Leu Glu Gly Gly Leu Leu Leu Leu  
20 25 30

5 Leu Leu Leu Val Thr Ala Ala Leu Val Ala Leu Gly Val Leu Tyr Ala  
35 40 45

Asp Arg Arg Gly Lys Gln Leu Pro Arg Leu Ala Ser Arg Leu Cys Phe  
50 55 60

10 Leu Gln Glu Glu Arg Thr Phe Val Lys Arg Lys Pro Arg Gly Ile Pro  
65 70 75 80

15 Glu Ala Gln Glu Val Ser Glu Val Cys Thr Thr Pro Gly Cys Val Ile  
85 90 95

Ala Ala Ala Arg Ile Leu Gln Asn Met Asp Pro Thr Thr Glu Pro Cys  
100 105 110

20 Asp Asp Phe Tyr Gln Phe Ala Cys Gly Gly Trp Leu Arg Arg His Val  
115 120 125

Ile Pro Glu Thr Asn Ser Arg Tyr Ser Ile Phe Asp Val Leu Arg Asp  
130 135 140

25 Glu Leu Glu Val Ile Leu Lys Ala Val Leu Glu Asn Ser Thr Ala Lys  
145 150 155 160

Asp Arg Pro Ala Val Glu Lys Ala Arg Thr Leu Tyr Arg Ser Cys Met  
165 170 175

30 Asn Gln Ser Val Ile Glu Lys Arg Gly Ser Gln Pro Leu Leu Asp Ile  
180 185 190

35 Leu Glu Val Val Gly Gly Trp Pro Val Ala Met Asp Arg Trp Asn Glu  
195 200 205

Thr Val Gly Leu Glu Trp Glu Leu Glu Arg Gln Leu Ala Leu Met Asn  
210 215 220

40 Ser Gln Phe Asn Arg Arg Val Leu Ile Asp Leu Phe Ile Trp Asn Asp  
225 230 235 240

Asp Gln Asn Ser Ser Arg His Ile Ile Tyr Ile Asp Gln Pro Thr Leu  
245 250 255

45 Gly Met Pro Ser Arg Glu Tyr Tyr Phe Asn Gly Gly Ser Asn Arg Lys  
260 265 270

50 Val Arg Glu Ala Tyr Leu Gln Phe Met Val Ser Val Ala Thr Leu Leu  
275 280 285

Arg Glu Asp Ala Asn Leu Pro Arg Asp Ser Cys Leu Val Gln Glu Asp  
290 295 300

55 Met Met Gln Val Leu Glu Leu Glu Thr Gln Leu Ala Lys Ala Thr Val  
305 310 315 320

Pro Gln Glu Glu Arg His Asp Val Ile Ala Leu Tyr His Arg Met Gly  
325 330 335

60 Leu Glu Glu Leu Gln Ser Gln Phe Gly Leu Lys Gly Phe Asn Trp Thr  
340 345 350

64

Leu Phe Ile Gln Thr Val Leu Ser Ser Val Lys Ile Lys Leu Leu Pro  
 355 360 365

5 Asp Glu Glu Val Val Val Tyr Gly Ile Pro Tyr Leu Gln Asn Leu Glu  
 370 375 380

Asn Ile Ile Asp Thr Tyr Ser Ala Arg Thr Ile Gln Asn Tyr Leu Val  
 385 390 395 400

10 Trp Arg Leu Val Leu Asp Arg Ile Gly Ser Leu Ser Gln Arg Phe Lys  
 405 410 415

Asp Thr Arg Val Asn Tyr Arg Lys Ala Leu Phe Gly Thr Met Val Glu  
 420 425 430

15 Glu Val Arg Trp Arg Glu Cys Val Gly Tyr Val Asn Ser Asn Met Glu  
 435 440 445

20 Asn Ala Val Gly Ser Leu Tyr Val Arg Glu Ala Phe Pro Gly Asp Ser  
 450 455 460

Lys Ser Met Val Arg Glu Leu Ile Asp Lys Val Arg Thr Val Phe Val  
 465 470 475 480

25 Glu Thr Leu Asp Glu Leu Gly Trp Met Asp Glu Glu Ser Lys Lys Lys  
 485 490 495

Ala Gln Glu Lys Ala Met Ser Ile Arg Glu Gln Ile Gly His Pro Asp  
 500 505 510

30 Tyr Ile Leu Glu Glu Met Asn Arg Arg Leu Asp Glu Glu Tyr Ser Asn  
 515 520 525

Leu Asn Phe Ser Glu Asp Leu Tyr Phe Glu Asn Ser Leu Gln Asn Leu  
 530 535 540

35 Lys Val Gly Ala Gln Arg Ser Leu Arg Lys Leu Arg Glu Lys Val Asp  
 545 550 555 560

40 Pro Asn Leu Trp Ile Ile Gly Ala Ala Val Val Asn Ala Phe Tyr Ser  
 565 570 575

Pro Asn Arg Asn Gln Ile Val Phe Pro Ala Gly Ile Leu Gln Pro Pro  
 580 585 590

45 Phe Phe Ser Lys Glu Gln Pro Gln Ala Leu Asn Phe Gly Gly Ile Gly  
 595 600 605

50 Met Val Ile Gly His Glu Ile Thr His Gly Phe Asp Asp Asn Gly Arg  
 610 615 620

Asn Phe Asp Lys Asn Gly Asn Met Met Asp Trp Trp Ser Asn Phe Ser  
 625 630 635 640

55 Thr Gln His Phe Arg Glu Gln Ser Glu Cys Met Ile Tyr Gln Tyr Gly  
 645 650 655

Asn Tyr Ser Trp Asp Leu Ala Asp Glu Gln Asn Val Asn Gly Phe Asn  
 660 665 670

60 Thr Leu Gly Glu Asn Ile Ala Asp Asn Gly Gly Val Arg Gln Ala Tyr  
 675 680 685



65

Lys Ala Tyr Leu Lys Trp Met Ala Glu Gly Gly Lys Asp Gln Gln Leu  
690 695 700

5 Pro Gly Leu Asp Leu Thr His Glu Gln Leu Phe Phe Ile Asn Tyr Ala  
705 710 715 720

Gln Val Trp Cys Gly Ser Tyr Arg Pro Glu Phe Ala Ile Gln Ser Ile  
725 730 735

10 Lys Thr Asp Val His Ser Pro Leu Lys Tyr Arg Val Leu Gly Ser Leu  
740 745 750

Gln Asn Leu Ala Ala Phe Ala Asp Thr Phe His Cys Ala Arg Gly Thr  
755 760 765

15 Pro Met His Pro Lys Glu Arg Cys Arg Val Trp  
770 775

20 <210> 5  
<211> 2262  
<212> DNA  
<213> Homo Sapiens

25 <220>  
<221> CDS  
<222> (1)..(2259)

30 <400> 5  
atg ggg aag tcc gaa ggc cca gtg ggg atg gtg gag agc gcc ggc cgt 48  
Met Gly Lys Ser Glu Gly Pro Val Gly Met Val Glu Ser Ala Gly Arg  
1 5 10 15

35 gca ggg cag aag cgc ccg ggg ttc ctg gag ggg ggg ctg ctg ctg ctg 96  
Ala Gly Gln Lys Arg Pro Gly Phe Leu Glu Gly Gly Leu Leu Leu Leu  
20 25 30

40 ctg ctg ctg gtg acc gct gcc ctg gtg gcc ttg ggt gtc ctc tac gcc 144  
Leu Leu Leu Val Thr Ala Ala Leu Val Ala Leu Gly Val Leu Tyr Ala  
35 40 45

45 gac cgc aga ggg atc cca gag gcc caa gag gtg agc gag gtc tgc acc 192  
Asp Arg Arg Gly Ile Pro Glu Ala Gln Glu Val Ser Glu Val Cys Thr  
50 55 60

50 acc cct ggc tgc gtg ata gca gct gcc agg atc ctc cag aac atg gac 240  
Thr Pro Gly Cys Val Ile Ala Ala Ala Arg Ile Leu Gln Asn Met Asp  
65 70 75 80

55 ccg acc acg gaa ccg tgt gac gac ttc tac cag ttt gca tgc gga ggc 288  
Pro Thr Thr Glu Pro Cys Asp Asp Phe Tyr Gln Phe Ala Cys Gly Gly  
85 90 95

60 tgg ctg cgg cgc cac gtg atc cct gag acc aac tca aga tac agc atc 336  
Trp Leu Arg Arg His Val Ile Pro Glu Thr Asn Ser Arg Tyr Ser Ile  
100 105 110

65 ttt gac gtc ctc cgc gac gag ctg gag gtc atc ctc aaa gcg gtg ctg 384  
Phe Asp Val Leu Arg Asp Glu Leu Glu Val Ile Leu Lys Ala Val Leu  
115 120 125

## 66

	gag aat tgc aot gcc aag gac cgg ccg gct gtg gag aag gcc agg acg	432
	Glu Asn Ser Thr Ala Lys Asp Arg Pro Ala Val Glu Lys Ala Arg Thr	
	130 135 140	
5	ctg tac cgc tcc tgc atg aac cag agt gtg ata gag aag cga ggc tct	480
	Leu Tyr Arg Ser Cys Met Asn Gln Ser Val Ile Glu Lys Arg Gly Ser	
	145 150 155 160	
10	cag ccc ctg ctg gac atc ttg gag gtg gtg gga ggc tgg ccg gtg gcg	528
	Gln Pro Leu Leu Asp Ile Leu Glu Val Val Gly Gly Trp Pro Val Ala	
	165 170 175	
15	atg gac agg tgg aac gag acc gta gga ctc gag tgg gag ctg gag cgg	576
	Met Asp Arg Trp Asn Glu Thr Val Gly Leu Glu Trp Glu Leu Glu Arg	
	180 185 190	
20	cag ctg gcg ctg atg aac tca cag ttc aac agg cgc gtc ctc atc gac	624
	Gln Leu Ala Leu Met Asn Ser Gln Phe Asn Arg Arg Val Leu Ile Asp	
	195 200 205	
25	ctc ttc atc tgg aac gac gac cag aac tcc agc cgg cac atc atc tac	672
	Leu Phe Ile Trp Asn Asp Asp Gln Asn Ser Ser Arg His Ile Ile Tyr	
	210 215 220	
30	ata gac cag ccc acc ttg ggc atg ccc tcc cga gag tac tac ttc aac	720
	Ile Asp Gln Pro Thr Leu Gly Met Pro Ser Arg Glu Tyr Tyr Phe Asn	
	225 230 235 240	
35	ggc ggc agc aac cgg aag gtg cgg gaa gcc tac ctg cag ttc atg gtg	768
	Gly Gly Ser Asn Arg Lys Val Arg Glu Ala Tyr Leu Gln Phe Met Val	
	245 250 255	
40	tca gtg gcc acg ttg ctg cgg gag gat gca aac ctg ccc agg gac agc	816
	Ser Val Ala Thr Leu Leu Arg Glu Asp Ala Asn Leu Pro Arg Asp Ser	
	260 265 270	
45	tgc ctg gtg cag gag gac atg atg cag gtg ctg gag ctg gag aca cag	864
	Cys Leu Val Gln Glu Asp Met Met Gln Val Leu Glu Leu Glu Thr Gln	
	275 280 285	
50	ctg gcc aag gcc acg gta ccc cag gag gag aga cac gac gtc atc gcc	912
	Leu Ala Lys Ala Thr Val Pro Gln Glu Glu Arg His Asp Val Ile Ala	
	290 295 300	
55	ttg tac cac cgg atg gga ctg gag gag ctg caa agc cag ttt ggc ctg	960
	Leu Tyr His Arg Met Gly Leu Glu Glu Leu Gln Ser Gln Phe Gly Leu	
	305 310 315 320	
60	aag gga ttt aac tgg act ctg ttc ata caa act gtg cta tcc tct gtc	1008
	Lys Gly Phe Asn Trp Thr Leu Phe Ile Gln Thr Val Leu Ser Ser Val	
	325 330 335	
65	aaa atc aag ctg ctg cca gat gag gaa gtg gtg gtc tat ggc atc ccc	1056
	Lys Ile Lys Leu Leu Pro Asp Glu Glu Val Val Val Tyr Gly Ile Pro	
	340 345 350	
70	tac ctg cag aac ctt gaa aac atc atc gac acc tac tca gcc agg acc	1104
	Tyr Leu Gln Asn Leu Glu Asn Ile Ile Asp Thr Tyr Ser Ala Arg Thr	
	355 360 365	
75	ata cag aac tac ctg gtc tgg cgc ctg gtg ctg gac cgc att ggt agc	1152
	Ile Gln Asn Tyr Leu Val Trp Arg Leu Val Leu Asp Arg Ile Gly Ser	
	370 375 380	

67

	cta	agc	cag	aga	ttc	aag	gac	aca	cga	gtg	aac	tac	cgc	aag	gcg	ctg	1200
	Leu	Ser	Gln	Arg	Phe	Lys	Asp	Thr	Arg	Val	Asn	Tyr	Arg	Lys	Ala	Leu	
	385					390					395					400	
5	ttt	ggc	aca	atg	gtg	gag	gag	gtg	cgc	tgg	cgt	gaa	tgt	gtg	ggc	tac	1248
	Phe	Gly	Thr	Met	Val	Glu	Glu	Val	Arg	Trp	Arg	Glu	Cys	Val	Gly	Tyr	
					405					410					415		
10	gtc	aac	agc	aac	atg	gag	aac	gcc	gtg	ggc	tcc	ctc	tac	gtc	agg	gag	1296
	Val	Asn	Ser	Asn	Met	Glu	Asn	Ala	Val	Gly	Ser	Leu	Tyr	Val	Arg	Glu	
				420					425					430			
15	gcg	ttc	cct	gga	gac	agc	aag	agc	atg	gtc	aga	gaa	ctc	att	gac	aag	1344
	Ala	Phe	Pro	Gly	Asp	Ser	Lys	Ser	Met	Val	Arg	Glu	Leu	Ile	Asp	Lys	
			435					440					445				
20	gtg	cgg	aca	gtg	ttt	gtg	gag	acg	ctg	gac	gag	ctg	ggc	tgg	atg	gac	1392
	Val	Arg	Thr	Val	Phe	Val	Glu	Thr	Leu	Asp	Glu	Leu	Gly	Trp	Met	Asp	
		450				455						460					
25	gag	gag	tcc	aag	aag	aag	gag	cag	gag	aag	gcc	atg	agc	atc	cgg	gag	1440
	Glu	Glu	Ser	Lys	Lys	Lys	Ala	Gln	Glu	Lys	Ala	Met	Ser	Ile	Arg	Glu	
	465				470					475						480	
30	cag	atc	ggg	cac	cct	gac	tac	atc	ctg	gag	gag	atg	aac	agg	cgc	ctg	1488
	Gln	Ile	Gly	His	Pro	Asp	Tyr	Ile	Leu	Glu	Glu	Met	Asn	Arg	Arg	Leu	
				485					490						495		
35	gac	gag	gag	tac	tcc	aat	ctg	aac	ttc	tca	gag	gac	ctg	tac	ttt	gag	1536
	Asp	Glu	Glu	Tyr	Ser	Asn	Leu	Asn	Phe	Ser	Glu	Asp	Leu	Tyr	Phe	Glu	
				500				505						510			
40	aac	agt	ctg	cag	aac	ctc	aag	gtg	ggc	gcc	cag	cgg	agc	ctc	agg	aag	1584
	Asn	Ser	Leu	Gln	Asn	Leu	Lys	Val	Gly	Ala	Gln	Arg	Ser	Leu	Arg	Lys	
			515				520						525				
45	ctt	cgg	gaa	aag	gtg	gac	cca	aat	ctc	tgg	atc	atc	ggg	ggc	ggc	gtg	1632
	Leu	Arg	Glu	Lys	Val	Asp	Pro	Asn	Leu	Trp	Ile	Ile	Gly	Ala	Ala	Val	
		530				535						540					
50	gtc	aat	gag	ttc	tac	tcc	cca	aac	cga	aac	cag	att	gta	ttc	cct	gcc	1680
	Val	Asn	Ala	Phe	Tyr	Ser	Pro	Asn	Arg	Asn	Gln	Ile	Val	Phe	Pro	Ala	
	545					550				555						560	
55	ggg	atc	ctc	cag	ccc	ccc	ttc	ttc	agc	aag	gag	cag	cca	cag	gcc	ttg	1728
	Gly	Ile	Leu	Gln	Pro	Pro	Phe	Phe	Ser	Lys	Glu	Gln	Pro	Gln	Ala	Leu	
				565					570						575		
60	aac	ttt	gga	ggc	att	ggg	atg	gtg	atc	ggg	cac	gag	atc	acg	cac	ggc	1776
	Asn	Phe	Gly	Gly	Ile	Gly	Met	Val	Ile	Gly	His	Glu	Ile	Thr	His	Gly	
				580				585						590			
65	ttt	gac	gac	aat	ggc	cgg	aac	ttc	gac	aag	aat	ggc	aac	atg	atg	gat	1824
	Phe	Asp	Asp	Asn	Gly	Arg	Asn	Phe	Asp	Lys	Asn	Gly	Asn	Met	Met	Asp	
			595				600						605				
70	tgg	tgg	agt	aac	ttc	tcc	acc	cag	cac	ttc	cgg	gag	cag	tca	gag	tgc	1872
	Trp	Trp	Ser	Asn	Phe	Ser	Thr	Gln	His	Phe	Arg	Glu	Gln	Ser	Glu	Cys	
		610					615					620					

68

atg atc tac cag tac ggc aac tac tcc tgg gac ctg gca gac gaa cag 1920  
 Met Ile Tyr Gln Tyr Gly Asn Tyr Ser Trp Asp Leu Ala Asp Glu Gln  
 625 630 635 640

5 aac gtg aac gga ttc aac acc ctt ggg gaa aac att gct gac aac gga 1968  
 Asn Val Asn Gly Phe Asn Thr Leu Gly Glu Asn Ile Ala Asp Asn Gly  
 645 650 655

10 ggg gtg cgg caa gcc tat aag gcc tac ctc aag tgg atg gca gag ggt 2016  
 Gly Val Arg Gln Ala Tyr Lys Ala Tyr Leu Lys Trp Met Ala Glu Gly  
 660 665 670

15 ggc aag gac cag cag ctg ccc ggc ctg gat ctc acc cat gag cag ctc 2064  
 Gly Lys Asp Gln Gln Leu Pro Gly Leu Asp Leu Thr His Glu Gln Leu  
 675 680 685

20 ttc ttc atc aac tac gcc cag gtg tgg tgc ggg tcc tac cgg ccc gag 2112  
 Phe Phe Ile Asn Tyr Ala Gln Val Trp Cys Gly Ser Tyr Arg Pro Glu  
 690 695 700

ttc gcc atc caa tcc atc aag aca gac gtc cac agt ccc ctg aag tac 2160  
 Phe Ala Ile Gln Ser Ile Lys Thr Asp Val His Ser Pro Leu Lys Tyr  
 705 710 715 720

25 agg gta ctg ggg tgg ctg cag aac ctg gcc gcc ttc gca gac acg ttc 2208  
 Arg Val Leu Gly Ser Leu Gln Asn Leu Ala Ala Phe Ala Asp Thr Phe  
 725 730 735

30 cac tgt gcc cgg ggc acc ccc atg cac ccc aag gag cga tgc cgc gtg 2256  
 His Cys Ala Arg Gly Thr Pro Met His Pro Lys Glu Arg Cys Arg Val  
 740 745 750

tgg tag 2262  
 Trp

35

<210> 6  
 <211> 753  
 <212> PRT  
 40 <213> Homo Sapiens

<400> 6  
 Met Gly Lys Ser Glu Gly Pro Val Gly Met Val Glu Ser Ala Gly Arg  
 1 5 10 15

45 Ala Gly Gln Lys Arg Pro Gly Phe Leu Glu Gly Gly Leu Leu Leu Leu  
 20 25 30

50 Leu Leu Leu Val Thr Ala Ala Leu Val Ala Leu Gly Val Leu Tyr Ala  
 35 40 45

Asp Arg Arg Gly Ile Pro Glu Ala Gln Glu Val Ser Glu Val Cys Thr  
 50 55 60

55 Thr Pro Gly Cys Val Ile Ala Ala Ala Arg Ile Leu Gln Asn Met Asp  
 65 70 75 80

Pro Thr Thr Glu Pro Cys Asp Asp Phe Tyr Gln Phe Ala Cys Gly Gly  
 85 90 95

60 Trp Leu Arg Arg His Val Ile Pro Glu Thr Asn Ser Arg Tyr Ser Ile  
 100 105 110

89

	Phe	Asp	Val	Leu	Arg	Asp	Glu	Leu	Glu	Val	Ile	Leu	Lys	Ala	Val	Leu	
			115					120					125				
5	Glu	Asn	Ser	Thr	Ala	Lys	Asp	Arg	Pro	Ala	Val	Glu	Lys	Ala	Arg	Thr	
			130				135					140					
	Leu	Tyr	Arg	Ser	Cys	Met	Asn	Gln	Ser	Val	Ile	Glu	Lys	Arg	Gly	Ser	
						150					155					160	
10	Gln	Pro	Leu	Leu	Asp	Ile	Leu	Glu	Val	Val	Gly	Gly	Trp	Pro	Val	Ala	
					165					170					175		
	Met	Asp	Arg	Trp	Asn	Glu	Thr	Val	Gly	Leu	Glu	Trp	Glu	Leu	Glu	Arg	
				180					185					190			
15	Gln	Leu	Ala	Leu	Met	Asn	Ser	Gln	Phe	Asn	Arg	Arg	Val	Leu	Ile	Asp	
			195					200					205				
20	Leu	Phe	Ile	Trp	Asn	Asp	Asp	Gln	Asn	Ser	Ser	Arg	His	Ile	Ile	Tyr	
		210				215						220					
	Ile	Asp	Gln	Pro	Thr	Leu	Gly	Met	Pro	Ser	Arg	Glu	Tyr	Tyr	Phe	Asn	
		225				230					235					240	
25	Gly	Gly	Ser	Asn	Arg	Lys	Val	Arg	Glu	Ala	Tyr	Leu	Gln	Phe	Met	Val	
				245						250					255		
30	Ser	Val	Ala	Thr	Leu	Leu	Arg	Glu	Asp	Ala	Asn	Leu	Pro	Arg	Asp	Ser	
				260					265					270			
	Cys	Leu	Val	Gln	Glu	Asp	Met	Met	Gln	Val	Leu	Glu	Leu	Glu	Thr	Gln	
			275					280					285				
35	Leu	Ala	Lys	Ala	Thr	Val	Pro	Gln	Glu	Glu	Arg	His	Asp	Val	Ile	Ala	
		290					295					300					
	Leu	Tyr	His	Arg	Met	Gly	Leu	Glu	Glu	Leu	Gln	Ser	Gln	Phe	Gly	Leu	
		305				310					315					320	
40	Lys	Gly	Phe	Asn	Trp	Thr	Leu	Phe	Ile	Gln	Thr	Val	Leu	Ser	Ser	Val	
				325						330					335		
45	Lys	Ile	Lys	Leu	Leu	Pro	Asp	Glu	Glu	Val	Val	Val	Tyr	Gly	Ile	Pro	
				340					345					350			
	Tyr	Leu	Gln	Asn	Leu	Glu	Asn	Ile	Ile	Asp	Thr	Tyr	Ser	Ala	Arg	Thr	
			355					360					365				
50	Ile	Gln	Asn	Tyr	Leu	Val	Trp	Arg	Leu	Val	Leu	Asp	Arg	Ile	Gly	Ser	
		370					375					380					
	Leu	Ser	Gln	Arg	Phe	Lys	Asp	Thr	Arg	Val	Asn	Tyr	Arg	Lys	Ala	Leu	
		385				390					395					400	
55	Phe	Gly	Thr	Met	Val	Glu	Glu	Val	Arg	Trp	Arg	Glu	Cys	Val	Gly	Tyr	
				405						410					415		
60	Val	Asn	Ser	Asn	Met	Glu	Asn	Ala	Val	Gly	Ser	Leu	Tyr	Val	Arg	Glu	
				420					425					430			
	Ala	Phe	Pro	Gly	Asp	Ser	Lys	Ser	Met	Val	Arg	Glu	Leu	Ile	Asp	Lys	
			435					440					445				

70

Val Arg Thr Val Phe Val Glu Thr Leu Asp Glu Leu Gly Trp Met Asp  
 450 455 460

5 Glu Glu Ser Lys Lys Lys Ala Gln Glu Lys Ala Met Ser Ile Arg Glu  
 465 470 475 480

Gln Ile Gly His Pro Asp Tyr Ile Leu Glu Glu Met Asn Arg Arg Leu  
 485 490 495

10 Asp Glu Glu Tyr Ser Asn Leu Asn Phe Ser Glu Asp Leu Tyr Phe Glu  
 500 505 510

Asn Ser Leu Gln Asn Leu Lys Val Gly Ala Gln Arg Ser Leu Arg Lys  
 515 520 525

15 Leu Arg Glu Lys Val Asp Pro Asn Leu Trp Ile Ile Gly Ala Ala Val  
 530 535 540

20 Val Asn Ala Phe Tyr Ser Pro Asn Arg Asn Gln Ile Val Phe Pro Ala  
 545 550 555 560

Gly Ile Leu Gln Pro Pro Phe Phe Ser Lys Glu Gln Pro Gln Ala Leu  
 565 570 575

25 Asn Phe Gly Gly Ile Gly Met Val Ile Gly His Glu Ile Thr His Gly  
 580 585 590

Phe Asp Asp Asn Gly Arg Asn Phe Asp Lys Asn Gly Asn Met Met Asp  
 595 600 605

30 Trp Trp Ser Asn Phe Ser Thr Gln His Phe Arg Glu Gln Ser Glu Cys  
 610 615 620

35 Met Ile Tyr Gln Tyr Gly Asn Tyr Ser Trp Asp Leu Ala Asp Glu Gln  
 625 630 635 640

Asn Val Asn Gly Phe Asn Thr Leu Gly Glu Asn Ile Ala Asp Asn Gly  
 645 650 655

40 Gly Val Arg Gln Ala Tyr Lys Ala Tyr Leu Lys Trp Met Ala Glu Gly  
 660 665 670

Gly Lys Asp Gln Gln Leu Pro Gly Leu Asp Leu Thr His Glu Gln Leu  
 675 680 685

45 Phe Phe Ile Asn Tyr Ala Gln Val Trp Cys Gly Ser Tyr Arg Pro Glu  
 690 695 700

50 Phe Ala Ile Gln Ser Ile Lys Thr Asp Val His Ser Pro Leu Lys Tyr  
 705 710 715 720

Arg Val Leu Gly Ser Leu Gln Asn Leu Ala Ala Phe Ala Asp Thr Phe  
 725 730 735

55 His Cys Ala Arg Gly Thr Pro Met His Pro Lys Glu Arg Cys Arg Val  
 740 745 750

Trp

**Claims**

1. Use of a compound having combined, in particular concurrent, inhibitory activity

c) on neutral endopeptidase (NEP) and

5 d) on the metalloprotease IGS5 which is a polypeptide comprising an amino acid sequence which has at least 70% identity to one of the amino acid sequences selected from the group of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6;

or of a pharmaceutically acceptable salt or solvate or biolabile ester thereof, for the manufacture of a medicament (pharmaceutical composition) for treating a mammal, preferably a human,  
10 suffering from or being susceptible to a condition which can be alleviated or prevented by combined or concurrent inhibition of NEP and IGS5.

2. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to claim 1 for the manufacture of a medicament (pharmaceutical composition)  
15 for treating a mammal, preferably a human, suffering from or being susceptible to a disease or condition where big-ET-1 levels are elevated and which disease or condition can be alleviated or prevented by combined or concurrent inhibition of NEP and IGS5.

3. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to claim 1 for the manufacture of a medicament (pharmaceutical composition)  
20 for treating a mammal, preferably a human, suffering from or being susceptible to a disease or condition where ET-1 is significantly upregulated and which disease or condition can be alleviated or prevented by combined or concurrent inhibition of NEP and IGS5.

25 4. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to any of the claims 1 to 3 for the manufacture of a medicament (pharmaceutical composition) for treatment and/or prophylaxis of hypertension, including secondary forms of hypertension such as renal or pulmonary hypertension, heart failure, angina pectoris, arrhythmias, myocardial infarction, cardiac hypertrophy, cerebral ischemia, peripheral vascular disease,  
30 subarachnoidal hemorrhage, chronic obstructive pulmonary disease (COPD), asthma, renal disease, atherosclerosis, and pain in colorectal cancer or prostate cancer, in larger mammals, preferably in humans.

5. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to any of the claims 2 or 3 for the manufacture of a medicament  
35 (pharmaceutical composition) for treatment and/or prophylaxis of atherosclerosis in a larger mammal, preferably in a human.


6. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to any of the claims 2 or 3 for the manufacture of a medicament (pharmaceutical composition) for treatment and/or prophylaxis of hypertension, including secondary forms of hypertension such as renal or pulmonary hypertension, heart failure, angina pectoris, arrhythmias, myocardial infarction, cardiac hypertrophy, cerebral ischemia, peripheral vascular disease, subarachnoidal hemorrhage, chronic obstructive pulmonary disease (COPD), asthma, renal disease, atherosclerosis, and pain in colorectal cancer or prostate cancer, in larger mammals, preferably in humans.

7. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to any of the claims 1 to 6 characterized in that the metalloprotease IGS5 is a polypeptide having an amino acid sequence which has at least 70% identity to one of the amino acid sequences selected from the group of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6.

8. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to any of the claims 1 to 7 characterized in that the amino acid sequence comprised by or being the metalloprotease IGS5 has at least 95% identity to one of the amino acid sequences selected from the group of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6.



9. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to claim 8 characterized in that the amino acid sequence comprised by or being the metalloprotease IGS5 is identical to one of the amino acid sequences selected from the group of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6.

10. Use of a compound of formula I,



wherein

A stands for a group with formula II or III





73

in which formula II

$R^{1a}$  stands for a phenyl-lower-alkyl group which can be optionally substituted in the phenyl ring by lower alkyl, lower alkoxy or halogen, or for a naphthyl-lower-alkyl group,

$R^{2a}$  means hydrogen or a group forming a biolabile ester; and

5 in which formula III

$R^{1b}$  is hydrogen or a group forming a biolabile phosphonic acid ester,

$R^{2b}$  is hydrogen or a group forming a biolabile phosphonic acid ester;

and wherein

10:  $R^3$  means hydrogen or a group forming a biolabile carboxylic acid ester;

or a pharmaceutically acceptable salt or solvate or biolabile ester thereof for the manufacture of a medicament (pharmaceutical composition) for treating a larger mammal, preferably a human, suffering from or being susceptible to a disease or condition which can be alleviated or prevented


15 by combined or concurrent inhibition

a) of neutral endopeptidase (NEP) and

b) of the metalloprotease IGS5 which is a polypeptide comprising an amino acid sequence which has at least 70% identity to one of the amino acid sequences selected from the group of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6.

20

11. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to claim 10 characterised in that the compound of formula I is selected from the subgroup of compounds having a structure of formula Ia,



25

Ia


wherein

$R^{1a}$  stands for a phenyl-lower-alkyl group which can be optionally substituted in the phenyl ring by lower alkyl, lower alkoxy or halogen, or for a naphthyl-lower-alkyl group,

$R^{2a}$  means hydrogen or a group forming a biolabile ester, and

30  $R^3$  means hydrogen or a group forming a biolabile ester, and physiologically acceptable salts of acids or solvates thereof.

12. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to claim 10 characterised in that the compound of formula I is selected from the subgroup of compounds having a structure of formula Ib,



Ib

wherein

R<sup>1b</sup> is hydrogen or a group forming a biolabile phosphonic acid ester,

R<sup>2b</sup> is hydrogen or a group forming a biolabile phosphonic acid ester, and

R<sup>3</sup> is hydrogen or a group forming a biolabile carboxylic acid ester,

10 and physiologically acceptable salts of acids or solvates thereof.

13. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to any of the claims 10 to 12 for the manufacture of a medicament (pharmaceutical composition) for treating a mammal, preferably a human, suffering from or being  
15 susceptible to a disease or condition where big-ET-1 levels are elevated and which disease (condition) can be alleviated or prevented by combined or concurrent inhibition of NEP and IGS5.

14. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to any of the claims 10 to 12 for the manufacture of a medicament  
20 (pharmaceutical composition) for treating a mammal, preferably a human, suffering from or being susceptible to a disease (condition) where ET-1 is significantly upregulated and which disease (condition) can be alleviated or prevented by combined or concurrent inhibition of NEP and IGS5.

15. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester  
25 thereof according to any of the claims 10 to 14 for the manufacture of a medicament (pharmaceutical composition) for treatment and/or prophylaxis of hypertension, including secondary forms of hypertension such as renal or pulmonary hypertension, heart failure, angina pectoris, arrhythmias, myocardial infarction, cardiac hypertrophy, cerebral ischemia, peripheral vascular disease, subarachnoidal hemorrhage, chronic obstructive pulmonary disease (COPD),  
30 asthma, renal disease, atherosclerosis, and pain in colorectal cancer or prostate cancer, in larger mammals, preferably in humans.

16. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to any of the claims 13 or 14 for the manufacture of a medicament (pharmaceutical composition) for treatment and/or prophylaxis of atherosclerosis in a larger mammal, preferably in a human.

5

17. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to any of the claims 13 or 14 for the manufacture of a medicament (pharmaceutical composition) for treatment and/or prophylaxis of hypertension, including secondary forms of hypertension such as renal or pulmonary hypertension, heart failure, angina pectoris, arrhythmias, myocardial infarction, cardiac hypertrophy, cerebral ischemia, peripheral vascular disease, subarachnoidal hemorrhage, chronic obstructive pulmonary disease (COPD), asthma, renal disease, atherosclerosis, and pain in colorectal cancer or prostate cancer, in mammals, preferably in humans.

10

15

18. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to any of the claims 10 to 17 characterised in that the metalloprotease IGS5 is a polypeptide having an amino acid sequence which has at least 70% identity to one of the amino acid sequences selected from the group of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6.

20

19. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to any of the claims 10 to 18 characterised in that the amino acid sequence comprised by or being the metalloprotease IGS5 has at least 95% identity to one of the amino acid sequences selected from the group of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6.

25

20. Use of a compound or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to claim 19 characterised in that the amino acid sequence comprised by or being the metalloprotease IGS5 is identical to one of the amino acid sequences selected from the group of SEQ ID NO:2, SEQ ID NO:4 and SEQ ID NO:6.

30

21. Use of a first compound showing combined or concurrent NEP/IGS5 inhibitory activity or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to any of the claims 1 to 20 in combination with at least one additional compound selected from the group of other individual and/or combined metalloprotease inhibitors than the combined NEP/IGS5 inhibitors, said additional compound preferably being selected from the group of ACE inhibitors, selective ECE inhibitors, selective NEP inhibitors, dual NEP/ECE inhibitors, and mixed inhibitors of these metalloproteases, for the manufacture of a medicament (pharmaceutical composition) for combination treatment and/or combination prophylaxis of any of the diseases or conditions as referenced in any of the claims 1 to 20.

35

22. Use according to claim 21 of said first compound in combination with at least one of said additional compounds, characterized in that the first compound compound has a structure of formula I as referenced in one of the claims 10 to 12, preferably a structure of formula Ia as referenced in claim 11 or of formula Ib as referenced in claim 12.

23. Use according to any of the claims 21 to 22 of said first compound in combination with at least one of said additional compounds, characterized in that the combination is co-effective, preferably synergistically effective.

10

24. Pharmaceutical composition, comprising co-effective, preferably synergistically effective, amounts of:

a first compound showing combined or concurrent NEP/IGS5 inhibitory activity or a pharmaceutically acceptable salt or solvate or biolabile ester thereof according to any of the claims 1 to 20; and of at least one additional compound selected from the group of other individual and/or combined metalloprotease inhibitors than the combined NEP/IGS5 inhibitors, said additional compound preferably being selected from the group of ACE inhibitors, selective ECE inhibitors, selective NEP inhibitors, dual NEP/ECE inhibitors, and mixed inhibitors of these metalloproteases, for combination treatment and/or combination prophylaxis of any of the diseases or conditions as referenced in any of the claims 1 to 20.

25. Pharmaceutical composition according to claim 24, comprising co-effective, preferably synergistically effective, amounts of said first compound and of at least one of said additional compounds characterized in that the first compound compound has a structure of formula I as referenced in one of the claims 10 to 12, preferably a structure of formula Ia as referenced in claim 11 or of formula Ib as referenced in claim 12.

**Abstract**

This invention relates to the use of a compound having combined, in particular concurrent, inhibitory activity on neutral endopeptidase (NEP) and on a novel metalloprotease designated IGS5, or of a pharmaceutically acceptable salt or solvate or biolabile ester thereof, for the manufacture of a medicament (pharmaceutical composition) for treating a larger mammal, preferably a human, suffering from or being susceptible to a disease or condition which can be alleviated or prevented by combined or concurrent inhibition of NEP and IGS5. In a particular aspect the present invention pertains to the use of said compounds with combined or concurrent NEP/IGS5 inhibitory activity for treating a larger mammal, preferably a human, suffering from or being susceptible to a disease or condition where big-ET-1 levels are elevated and which disease or condition can be alleviated or prevented by combined or concurrent inhibition of NEP and IGS5. In a further particular aspect the present invention pertains to the use of said compounds with combined or concurrent NEP/IGS5 inhibitory activity for treating a larger mammal, preferably a human, suffering from or being susceptible to a disease or condition where ET-1 is significantly upregulated and which disease or condition can be alleviated or prevented by combined or concurrent inhibition of NEP and IGS5. In the present invention said compounds with combined or concurrent NEP/IGS5-inhibitory activity preferably are used for the treatment and/or prophylaxis of hypertension, including secondary forms of hypertension such as renal or pulmonary hypertension, heart failure, angina pectoris, arrhythmias, myocardial infarction, cardiac hypertrophy, cerebral ischemia, peripheral vascular disease, subarachnoidal hemorrhage, chronic obstructive pulmonary disease (COPD), asthma, renal disease, atherosclerosis, and pain in colorectal cancer or prostate cancer, in mammals, preferably in humans, and more preferably in a patient sub-population suffering from or being susceptible to a disease or condition which can be alleviated or prevented by combined or concurrent inhibition of NEP and IGS5. Furthermore, it may be beneficial to additionally combine the said compounds showing combined or concurrent NEP/IGS5 inhibitory activity with other individual and/or combined metalloprotease inhibitors than the NEP/IGS5 inhibitors, e.g. with separate ACE- and/or ECE- and/or NEP-inhibitors and/or mixed inhibitors of these metalloproteases.

